

Integrals over Grassmannians and Random permutations

M. Adler* P. van Moerbeke†

October 8, 2001

Abstract

In testing the independence of two Gaussian populations, one computes the distribution of the sample canonical correlation coefficients, given that the actual correlation is zero. The “Laplace transform” of this distribution is not only an integral over the Grassmannian of p -dimensional planes in complex n -space, but is also related to a generalized hypergeometric function. Such integrals are solutions of Painlevé-like equations. They also have expansions, related to random words of length ℓ formed with an alphabet of p letters. Given that each letter appears in the word, the maximal length of the disjoint union of p increasing subsequences of the word clearly equals ℓ . But the maximal length of the disjoint union of $p - 1$ increasing subsequences leads to a non-trivial distribution. It is precisely this probability which appears in the expansion above.

Contents

1	Integrals over Grassmannians	11
----------	-------------------------------------	-----------

*Department of Mathematics, Brandeis University, Waltham, Mass 02454, USA. E-mail: adler@math.brandeis.edu. The support of a National Science Foundation grant # DMS-98-4-50790 is gratefully acknowledged.

†Department of Mathematics, Université de Louvain, 1348 Louvain-la-Neuve, Belgium and Brandeis University, Waltham, Mass 02454, USA. E-mail: vanmoerbeke@geom.ucl.ac.be and @math.brandeis.edu. The support of a National Science Foundation grant # DMS-98-4-50790, a Nato, a FNRS and a Francqui Foundation grant is gratefully acknowledged.

2	Jack polynomials	23
2.1	Young diagrams and Schur polynomials	23
2.2	Some useful formulae on hook length	25
2.3	Jack polynomials	28
3	Probability measures on partitions	32
3.1	Probability measure on the set $\mathbb{Y}$ of all partitions	32
3.2	Probability measures on the set $\mathbb{Y}_\ell$ of partitions of ℓ and random words	34
4	Expressing integrals as mathematical expectation on partitions and on random words	36
4.1	Expressing an integral on $Gr(p, \mathbb{F}^n)$ as a mathematical expectation on partitions	36
4.2	Expressing an integral over $Gr(p, \mathbb{C}^n)$ as a mathematical expectation on partitions and random words	38
5	Testing Statistical Independence of Gaussian Populations	42
5.1	The Wishart distribution	42
5.2	The canonical correlation coefficients	43
5.3	Distribution of the sample canonical correlations	44
6	Differential equations for the Grassmannian integrals and the hypergeometric functions	48
6.1	Differential equations for the Grassmannian integral	48
6.2	Hypergeometric functions, KP hierarchy and Painlevé equations	55
7	Differential equations for the Wishart and canonical correlations distributions	58
7.1	Laguerre probability on the tangent space to $Gr(p, \mathbb{F}^n)$ and the Wishart distribution	59
7.2	Jacobi probability on the tangent space to $Gr(p, \mathbb{F}^n)$ and the sample canonical correlation distribution	61
8	Appendix: The Pfaff-KP hierarchy and Virasoro constraints	63

Random words, longest increasing sequences and mean hook lengths

Consider the set of words

$$\pi \in S_\ell^p := \{\text{words } \pi \text{ of length } \ell, \text{ built from an alphabet } \{1, \dots, p\}\},$$

with the uniform probability distribution

$$P^{\ell,p}(\pi) = \frac{1}{p^\ell}. \quad (0.0.1)$$

The RSK correspondence (see section 2.1) between words and pairs of semi-standard and standard tableaux induces a probability measure on partitions

$$\lambda \in \mathbb{Y}_\ell = \{\text{partitions } \lambda \in \mathbb{Y} \text{ of weight } |\lambda| = \ell\}, \quad (0.0.2)$$

given by

$$P^{\ell,p}(\lambda) = \frac{f^\lambda s_\lambda(1^p)}{p^\ell}, \quad |\lambda| = \ell, \quad (0.0.3)$$

where s_λ is the Schur polynomial associated with the partition λ ,

$$1^p = (\overbrace{1, \dots, 1}^p, 0, 0, \dots) \quad \text{and} \quad f^\lambda = \#\{\text{standard tableaux of shape } \lambda\}, \quad (0.0.4)$$

with¹

$$(\text{support } P^{\ell,p}) \subseteq \mathbb{Y}_\ell^{(p)} := \{\lambda \in \mathbb{Y}_\ell, \text{ such that } \lambda_1^\top \leq p\}.$$

A subsequence σ of the word π is *weakly k -increasing*, if it can be written as

$$\sigma = \sigma_1 \cup \sigma_2 \cup \dots \cup \sigma_k, \quad (0.0.5)$$

where σ_i are disjoint weakly increasing subsequences of the word π , i.e., possibly with repetitions. The length of the longest increasing/decreasing subsequences is closely related to the shape of the associated partition, via the RSK correspondence :

$$\begin{aligned} d_1(\pi) &= \left\{ \begin{array}{l} \text{length of the longest } \textit{strictly} \\ \text{decreasing subsequence of } \pi \end{array} \right\} = \lambda_1^\top \\ i_k(\pi) &= \left\{ \begin{array}{l} \text{length of the longest } \textit{weakly} \\ k\text{-increasing subsequence of } \pi \end{array} \right\} = \lambda_1 + \dots + \lambda_k \end{aligned} \quad (0.0.6)$$

¹ $\lambda^\top$ is the dual partition, i.e., obtained by flipping the Young diagram λ about its diagonal. So, $\lambda_1^\top$ is the length of the first column of λ .


For integer $0 \leq p < n$, consider the fixed rectangular Young diagram of width $n - p$,

$$\mu = (n - p)^p := (\overbrace{n - p, n - p, \dots, n - p}^p). \quad (0.0.7)$$

Consider a word $\pi \in S_\ell^p$. Then the statement $d_1(\pi) = p$ implies, in particular, that all letters of the alphabet $\{1, \dots, p\}$ are represented in π ; then automatically $i_p(\pi) = \ell$. The theorem below deals with the first non-trivial quantity $i_{p-1}(\pi)$, given that $d_1(\pi) = p$. Using the standard notation, defined for a general parameter $\beta > 0$,

$$(a)_\lambda^{(1/\beta)} := \prod_i (a + \beta(1 - i))_{\lambda_i}, \text{ with } (x)_n := x(x+1) \dots (x+n-1), \ x_0 = 1, \quad (0.0.8)$$

we now state Theorem 0.1, which will be established in section 3.2 (note that here symbol (0.0.8) is used for $\beta = 1$):



Theorem 0.1 *Given the probability (0.0.1) and (0.0.3), the following holds (h^κ denotes the product of hook lengths over all boxes of the partition κ):*

$(0 \leq p < n)$

$$\begin{aligned} P^{\ell,p}(\lambda \supseteq \mu) &= P^{\ell,p} \left(\pi \in S_\ell^p \mid \begin{array}{l} d_1(\pi) = p \text{ and} \\ i_{p-1}(\pi) \leq \ell - n + p \end{array} \right) \\ &= \frac{\ell!}{p^\ell} \prod_1^p \frac{(p-i)!}{(n-i)!} \sum_{\substack{\kappa \in \mathbb{Y}_{\ell-p}(n-p) \\ \kappa_1^\top \leq p}} \frac{1}{(h^\kappa)^2} \frac{((p)_\kappa)^2}{(n)_\kappa}. \end{aligned} \quad (0.0.9)$$

More generally, for fixed $p \leq q < n$, the mathematical expectation (with regard to $P^{\ell,p}$) of the hook length of λ , emanating from the vertical strip ν (of width $q - p$), equals:

$$E^{\ell,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^\lambda \right) = \frac{\ell!}{p^\ell} \prod_1^p \frac{(q-i)!}{(n-i)!} \sum_{\substack{\kappa \in \mathbb{Y}_{\ell-p}(n-p) \\ \kappa_1^\top \leq p}} \frac{1}{(h^\kappa)^2} \frac{(p)_\kappa (q)_\kappa}{(n)_\kappa}. \quad (0.0.10)$$

Generating function for the mathematical expectation of the hook length, integrals over Grassmannians and Painlevé V

Theorem 0.2 below involves an *integral over the Grassmannian*

$$Gr(p, \mathbb{C}^n) = \frac{U(n)}{U(p) \times U(n-p)} =: G/K \quad (0.0.11)$$

of p -dimensional planes in $\mathbb{C}^n$ through the origin and Haar measure $d\mu(Z)$ on $Gr(p, \mathbb{C}^n)$, expressed in the parametrizing coordinate Z of

$$\text{Affine } Gr(p, \mathbb{C}^n) = \left\{ \text{span} \left(\begin{array}{c} I_p \\ Z \end{array} \right) \mid Z := A_{21} A_{11}^{-1}, \ A \in G \right\}, \quad (0.0.12)$$

where $A \in G$ is represented in block form:

$$\begin{array}{ccc} \xleftarrow{p} & \xleftarrow{n-p} & \\ \left(\begin{array}{cc} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right) & \begin{array}{c} \updownarrow p \\ \updownarrow n-p \end{array} & \in G. \end{array} \quad (0.0.13)$$

Section 1 will be devoted to the geometry of $Gr(p, \mathbb{F}^n)$, where $\mathbb{F} = \mathbb{C}, \mathbb{R}$ or $\mathbb{H}$, and to the study of integrals over $Gr(p, \mathbb{F}^n)$.

We also need *Jack polynomials*, which are the unique symmetric functions orthogonal with respect to a certain α -dependent inner-product $\langle, \rangle_\alpha$, such that

$$\langle J_\lambda^{(\alpha)}, J_\mu^{(\alpha)} \rangle = \delta_{\mu\lambda} j_\lambda^{(\alpha)}, \quad (0.0.14)$$

with

$$j_\lambda^{(\alpha)} = \prod_{(i,j) \in \lambda} (\lambda_j^\top - i + \alpha(\lambda_i - j + 1)) (\lambda_j^\top - i + 1 + \alpha(\lambda_i - j)).$$

Facts about Jack polynomials relevant for this project will be discussed in section 2.3.

Finally, *generalized hypergeometric functions* ${}_2F_1^{(\alpha)}$ are defined by: ($p, q, n \in \mathbb{C}$, $x = (x_1, x_2, \dots)$)

$${}_2F_1^{(\alpha)}(p, q; n; x) := \sum_{\kappa \in \mathbb{Y}} \frac{(p)_\kappa^{(\alpha)} (q)_\kappa^{(\alpha)}}{(n)_\kappa^{(\alpha)}} \alpha^{|\kappa|} \frac{J_\kappa^{(\alpha)}(x)}{j_\kappa^{(\alpha)}}. \quad (0.0.15)$$

In particular,

$${}_2F_1^{(1)}(p, q; n; x) := \sum_{\kappa \in \mathbb{Y}} \frac{(p)_\kappa^{(1)} (q)_\kappa^{(1)}}{(n)_\kappa^{(1)} h^\kappa} s_\kappa(x). \quad (0.0.16)$$

The following theorem will be established in section 4.2; it is strongly motivated by certain integrals appearing in the context of testing statistical independence of Gaussian populations, as will be explained in the next paragraph.

Theorem 0.2 *For fixed $p \leq q \leq n/2$, the generating function for the mathematical expectation of the hook length (0.0.10) over a strip, with regard to the probability (0.0.3), is given by²*

²Remember (support $P^{\ell,p}$) $\subseteq \{\lambda \in \mathbb{Y}, \text{ such that } |\lambda| = \ell, \lambda_1^\top \leq p\}$.

$$\begin{aligned}
& \prod_{i=1}^p \frac{(n-i)!}{(q-i)!} x^{-p(n-p)} \sum_{\ell \geq p(n-p)} \frac{(px)^\ell}{\ell!} E^{\ell,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^\lambda \right) \\
&= (c_{n,q,p}^{(1)})^{-1} \int_{Gr(p, \mathbb{C}^n)} e^{x \operatorname{Tr}(I + Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-(q-p)} d\mu(Z) \\
&= {}_2F_1^{(1)}(p, q; n; y) \Big|_{\sum_\ell y_\ell^i = \delta_{1i} x} \\
&= \exp \int_0^x \frac{u(y) - p(n-p) + py}{y} dy \tag{0.0.17}
\end{aligned}$$

where $u(x)$ is the unique solution to the initial value problem:

$$\begin{cases} x^2 u''' + x u'' + 6x u'^2 - 4u u' + 4Q u' - 2Q' u + 2R = 0 & \text{(Painlevé V)} \\ \text{with } u(x) = p(n-p) - \frac{p(n-q)}{n}x + \dots + a_{n+1}x^{n+1} + O(x^{n+1}) + \dots, \text{ near } x = 0. \end{cases} \tag{0.0.18}$$

with a_{n+1} specified by the hypergeometric function in (0.0.17) and

$$\begin{aligned}
4Q &= -x^2 + 2(n + 2(p-q))x - (n-2p)^2 \\
2R &= p(p-q)(x + n - 2p). \tag{0.0.19}
\end{aligned}$$

Remark: The constant $c_{n,q,p}^{(1)}$ in (0.0.17) is the one below for $\beta = 1$:

$$c := c_{n,q,p}^{(\beta)} := \prod_{i=1}^p \frac{\Gamma(i\beta + 1) \Gamma(\beta(n-q-i+1)) \Gamma(\beta(q-i+1))}{\Gamma(\beta+1) \Gamma(\beta(n-i+1))}. \tag{0.0.20}$$

Testing Statistical Independence of Gaussian Populations

To summarize section 5, consider $p+q$ normally distributed random variables $(X_1, \dots, X_p)^\top$ and $(Y_1, \dots, Y_q)^\top$ ($p \leq q$) with mean zero and covariance matrix Σ . According to Hotelling ([24])(see also Muirhead [35], p106), $(X_1, \dots, X_p)^\top$ and $(Y_1, \dots, Y_q)^\top$ can be replaced by a linearly transform of the X 's and Y 's, so that the covariance matrix takes on the canonical form:

$$\Sigma = \text{cov} \begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} \overset{p}{\longleftrightarrow} \Sigma_{11} & \overset{q}{\longleftrightarrow} \Sigma_{12} \\ \Sigma_{12}^\top & \Sigma_{22} \end{pmatrix} \begin{matrix} \updownarrow p \\ \updownarrow q \end{matrix} \longrightarrow \Sigma_{can} = \begin{pmatrix} I_p & P \\ P^\top & I_q \end{pmatrix},$$

where

$$P = \begin{pmatrix} \rho_1 & & & & & \\ & \ddots & & O & & \\ & & \rho_k & & & \\ & & & \rho_{k+1} & & \\ & O & & & \ddots & \\ & & & & & \rho_p \end{pmatrix} \begin{matrix} \xleftarrow{q} \xrightarrow{q} \\ \\ \\ \\ \\ \end{matrix} \left| \begin{matrix} O \\ \\ \\ \\ \\ \end{matrix} \right| \begin{matrix} \updownarrow p, \\ \updownarrow p, \\ \updownarrow p, \\ \updownarrow p, \\ \updownarrow p, \\ \updownarrow p, \end{matrix} \quad k = \text{rank } \Sigma_{12},$$

is the $p \times q$ -matrix of canonical correlation coefficients:

$$1 \geq \rho_1 \geq \rho_2 \geq \dots \geq \rho_k > 0, \quad \rho_{k+1} = \dots = \rho_p = 0.$$

The n ($n \geq p + q$) independent samples $(x_{11}, \dots, x_{1p}, y_{11}, \dots, y_{1q})^\top, \dots, (x_{n1}, \dots, x_{np}, y_{n1}, \dots, y_{nq})^\top$, arising from observing $\begin{pmatrix} X \\ Y \end{pmatrix}$ lead to a matrix $\begin{pmatrix} x \\ y \end{pmatrix}$ of size $(p+q, n)$, having a normal distribution with correlation Σ . The roots $r_1^2, \dots, r_p^2$ (sample canonical correlation coefficients) of the equation

$$\det(xy^\top(yy^\top)^{-1}yx^\top - r^2xx^\top) = 0$$

are the estimators (maximum likelihood estimators) of the canonical correlation coefficients $\rho_1^2, \dots, \rho_p^2$.

In testing the *null hypothesis*, versus the *alternative hypothesis*,

$$H_0: \rho_1^2 = \dots = \rho_p^2 = 0 \quad \text{versus} \quad H_a: (\rho_1^2, \dots, \rho_p^2) \neq 0$$

one needs the joint density of the r_i^2 , given that $\rho_1^2 = \dots = \rho_p^2 = 0$; namely, up to a (q, p, n) -dependent normalizing constant, the density is given by

$$|\Delta_p(r^2)|^\beta \prod_{i=1}^p (r_i^2)^{\beta(q-p+1)-1} (1 - r_i^2)^{\beta(n-q-p+1)-1} dr_i^2 \quad (0.0.21)$$

for $\beta = 1/2$. This formula generalizes to the formula above, upon considering random variables $(X_1, \dots, X_p)^\top$ and $(Y_1, \dots, Y_q)^\top$, with values in the complex $\mathbb{C}$ ($\beta = 1$) and the quaternions $\mathbb{H}$ ($\beta = 2$).

Expectation of the ratio of Jack polynomials, integrals over Grassmannians, sample canonical correlation coefficients and PDE's

Consider now a Poissonized probability on partitions $\lambda \in \mathbb{Y}$, which depends on a parameter $x > 0$:

$$P_{x,p}(\lambda) = e^{-\beta p x} \frac{x^{|\lambda|} J_{\lambda}^{(1/\beta)}(1^p)}{j_{\lambda}^{(1/\beta)}} = e^{-\beta p x} \frac{(\beta p x)^{|\lambda|}}{|\lambda|!} P^{\ell,p}(\lambda), \quad \lambda \in \mathbb{Y}, \quad (0.0.22)$$

with $P^{\ell,p}$ generalizing probability measure (0.0.3), as we shall see in section 3.1,

$$P^{\ell,p}(\lambda) = \frac{J_{\lambda}^{(1/\beta)}(1^p)}{j_{\lambda}^{(1/\beta)}} \frac{|\lambda|!}{(\beta p)^{|\lambda|}} \quad \text{with } |\lambda| = \ell,$$

This probability has its support on $\lambda_1^{\top} \leq p$. Many of these probability distributions on partitions have been introduced and extensively studied by Borodin, Kerov, Okounkov, Olshanski and Vershik (see [10, 11, 12, 50, 31]) The following statement involves an integral suggested again by the statistical theory mentioned earlier.

Theorem 0.3 *For fixed $p \leq q \leq n/2$, the following holds ($\beta = 1/2, 1, 2$)*

$$\begin{aligned} I_{n,p,q}^{(\beta)} &= c e^{\beta p x} E_{x,p} \left(\frac{1}{\beta^{|\lambda|}} \frac{J_{\lambda}^{(1/\beta)}(1^q)}{J_{\lambda}^{(1/\beta)}(1^n)} \right) \\ &= \int_{Gr(p, \mathbb{F}^n)} e^{x \operatorname{Tr}(I + Z^{\dagger} Z)^{-1}} \det(Z^{\dagger} Z)^{-\beta(q-p)} d\mu(Z) \\ &= \int_{[0,1]^p} e^{x \sum_1^p z_i} |\Delta_p(z)|^{2\beta} \prod_1^p z_i^{\beta(q-p+1)-1} (1 - z_i)^{\beta(n-q-p+1)-1} dz_i \\ &= c {}_2F_1^{(1/\beta)}(\beta p, \beta q; \beta n; y) \Big|_{\sum_{\ell} y_{\ell}^i = \frac{x}{\beta} \delta_{1i}} \\ &= c \exp \int_0^x v(y) dy. \end{aligned} \quad (0.0.23)$$

where $c = c_{n,q,p}^{(\beta)}$ is as in (0.0.20) and where

- $d\mu(Z)$ is Haar measure on the space $Gr(p, \mathbb{F}^n)$ of p -planes in $\mathbb{F}^n$, where $\mathbb{F} := \mathbb{C}, \mathbb{R}$ or the quaternions $\mathbb{H}$.

- The integral over $[0, 1]^p$, appearing in (0.0.23), is the “Laplace transform” of the distribution of the sample canonical correlation coefficients (0.0.21). This integral is a H ankel determinant for $\beta = 1$ and a Pfaffian for $\beta = 1/2$ and 2.
- $v(y) := v_{n,p,q}^{(\beta)}(y)$ and $I_p := I_{n,p,q}^{(\beta)}$ satisfies the differential equation (define $\delta_1^\beta := 1$ for $\beta = 1$ and $:= 0$ otherwise):

$$4 \left(y^3 v''' + 6y^3 v'^2 + (1 + \delta_1^\beta)(2y^2 v'' + 4y^2 v v' + y v^2) \right) - y P_0 v' + P_1 v + P_2$$

$$= \begin{cases} 0, & \text{for } \beta = 1, \text{ (Painlev   V)} \\ \frac{3}{16} \frac{p(p-1)}{(p+1)(p+2)} y^3 \frac{I_{p-2} I_{p+2}}{I_p^2}, & \text{for } \beta = 1/2, \\ \frac{3}{16^2} \frac{p}{p+1} y^3 \frac{I_{p-1} I_{p+1}}{I_p^2}, & \text{for } \beta = 2, \end{cases}$$

with P_0 quadratic and P_1, P_2 linear polynomials in y , with coefficients depending on n and

$$r = pq, \quad s = n - 2p - 2q. \quad (0.0.24)$$

This statement will be established in section 4.1 and the differential equation part in section 6.1. As a by-product, we show incidentally that the multivariate hypergeometric function ${}_2F_1^{(1)}(p, q; n; y)$ expressed in the $it_i = \sum_{k \geq 1} y_k^i$ -variables are τ -functions for the KP-hierarchy; but also that the function ${}_2F_1^{(1)}(p, q; n; y)$ properly restricted is a solution of Painlev   V. For related questions, see Orlov and Sherbin ([37, 38]). Section 7 gives new differential equations for the spectrum of Wishart matrices and for the sample canonical correlations of Gaussian populations.

Acknowledgment: The authors thank Professors I. Gessel, S. Helgason, B. Lian, G. Schwarz, R. Stanley and C.-L. Terng for useful advice, especially regarding section 1.

1 Integrals over Grassmannians

Consider the Grassmannian $Gr(p, \mathbb{F}^n)$ of p -planes through the origin in $\mathbb{F}^n$, where $\mathbb{F} = \mathbb{C}, \mathbb{R}$ or $\mathbb{H}$ (= quaternions). Let G be the group of matrices A , with entries in $\mathbb{F}$, such that $A^{-1} = A^\dagger$, with³ $A^\dagger := \bar{A}^\top$. Matrices $A \in G$ will be represented as block matrices

$$\begin{array}{ccc} \xleftrightarrow{p} & \xleftrightarrow{n-p} & \\ \left(\begin{array}{cc} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right) & \begin{array}{c} \updownarrow p \\ \updownarrow n-p \end{array} & \in G. \end{array} \quad (1.0.1)$$

The main statement of this section is theorem 1.1, where it is assumed, without loss of generality, that $n \geq 2p$. The values of β are related to $\mathbb{C}, \mathbb{R}$ and $\mathbb{H}$, as follows:

$$\begin{aligned} Gr(p, \mathbb{C}^n) : & \quad \beta = 1 \\ Gr(p, \mathbb{R}^n) : & \quad \beta = 1/2 \\ Gr(p, \mathbb{H}^n) : & \quad \beta = 2 \end{aligned}$$

The geometry of the symmetric spaces G/K and $K \backslash G/K$ has been studied by Helgason [22, 23]. In his recent Princeton thesis, Dueñez ([16]) has systematically studied integrals over symmetric spaces. Explicit information on this subject is not readily available in the literature; therefore we explain the theory in the Grassmannian case and the useful aspects for our purposes.

Theorem 1.1 *Consider the two parametrization of $Gr(p, \mathbb{F}^n)$*

$$Affine\ Gr(p, \mathbb{F}^n) = \left\{ span \left(\begin{array}{c} I_p \\ Z \end{array} \right) \mid Z := Z(A) = A_{21}A_{11}^{-1},\ A \in G \right\}, \quad (1.0.2)$$

and its invariant measures $d\mu(Z)$. Then, for $p \leq q \leq n/2$, we have

$$\begin{aligned} & \int_{Gr(p, \mathbb{F}^n)} e^{x \operatorname{Tr}(I + Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-\beta(q-p)} d\mu(Z) \\ &= \int_{[0,1]^p} e^{x \sum_1^p z_i} |\Delta_p(z)|^{2\beta} \prod_1^p z_i^{\beta(q-p+1)-1} (1 - z_i)^{\beta(n-q-p+1)-1} dz_i. \end{aligned} \quad (1.0.3)$$

³Given $a_i \in \mathbb{R}$, we define for $a = a_0 + ia_1 \in \mathbb{C}$, $\bar{a} = a_0 - ia_1$ and for $a = a_0 + a_1e_1 + a_2e_2 + a_3e_3 \in \mathbb{H}$, $\bar{a} := a_0 - a_1e_1 - a_2e_2 - a_3e_3$.

This is the “Fourier transform” of the joint density of the sample canonical correlations $(z_1, \dots, z_p) = (r_1^2, \dots, r_p^2)$, for $\beta = 1/2, 1, 2$, for the real, complex and quaternionic cases, given that the canonical correlation coefficients are zero. (see (0.0.21) and section 5)

Consider the following block matrix

$$I_{p,q} = \begin{pmatrix} I_p & O \\ O & -I_q \end{pmatrix}.$$

Theorem 1.2 *An alternative description for $Gr(p, \mathbb{F}^n)$ is given by*

$$Gr(p, \mathbb{F}^n) \simeq \mathcal{S} := \left\{ M = AI_{p,n-p}A^{-1}I_{p,n-p} \mid A \in G \right\} \quad (1.0.4)$$

The $n \times n$ matrices M have a $n - 2p$ -dimensional eigenspace corresponding to the eigenvalue 1, so that M can be decomposed into $M = M_0 \otimes M_1$ (with M_1 corresponding to the 1-eigenspace), with $Tr(M - M_0) = n - 2p$. Then we have

$$\begin{aligned} & e^{px/2} \int_{\mathcal{S}} e^{xTr M_0/4} \det \left(\frac{I - M_0}{I + M_0} \right)^{\beta(p-q)} d\mu(M) \\ &= \int_{[0,1]^p} e^{x \sum_1^p z_i} |\Delta_p(z)|^{2\beta} \prod_1^p z_i^{\beta(q-p+1)-1} (1 - z_i)^{\beta(n-q-p+1)-1} dz_i. \end{aligned} \quad (1.0.5)$$

Theorem 1.3 *Considering the parametrization*

$$T_{Id}Gr(p, \mathbb{F}^n) = \{ Z \mid \text{arbitrary } (n-p) \times p \text{ matrix, with } Z_{ij} \in \mathbb{F} \}, \quad (1.0.6)$$

we have

$$\begin{aligned} & \int_{\substack{Z \in T_{Id}Gr(p, \mathbb{F}^n) \\ \text{with spectrum } (Z^\dagger Z) \in E}} e^{-x Tr Z^\dagger Z} \det(Z^\dagger Z)^{\beta p} d\mu(Z), \quad E \subset \mathbb{R}^+ \\ &= \int_{E^p} |\Delta_p(u)|^{2\beta} \prod_1^p e^{-xu_i} u_i^{\beta(n-p+1)-1} du_i. \end{aligned} \quad (1.0.7)$$

For $\beta = 1/2$ and $x = \frac{1}{2\lambda}$, this is an integral of the joint density (Wishart density) (see section 5 and Muirhead [35], p.107) of the eigenvalues $u_1, \dots, u_p$ of the matrix $A = Z^\dagger Z$, where Z is a $n \times p$ matrix ($p \leq n$), with Gaussian density centered at 0 and covariance λI_p , namely the density

$$c_{n,p}(2\pi\lambda)^{-np/2} e^{-\frac{1}{2\lambda} \text{Tr } Z^\dagger Z} \prod_{\substack{1 \leq i \leq n \\ 1 \leq j \leq p}} dz_{ij}.$$

Proposition 1.4 *Then*

$$\begin{aligned} Gr(p, \mathbb{C}^n) &= \frac{U(n)}{U(p) \times U(n-p)} =: G/K \\ Gr(p, \mathbb{R}^n) &= \frac{SO(n)}{SO(p) \times SO(n-p)} =: G/K \\ Gr(p, \mathbb{H}^n) &= \frac{Sp(n)}{Sp(p) \times Sp(n-p)} =: G/K, \end{aligned} \quad (1.0.8)$$

and the affine part can be parametrized as follows:

$$\text{Affine } Gr(p, \mathbb{F}^n) = \left\{ \text{span} \begin{pmatrix} I_p \\ Z \end{pmatrix} \mid Z = Z(A) = A_{21}A_{11}^{-1}, A \in G \right\}$$

and

$$K \backslash G/K = \left\{ \text{span} \begin{pmatrix} I_p \\ Z \end{pmatrix} \mid Z = \begin{pmatrix} -\tan \theta_1 & & O \\ & \ddots & \\ O & & -\tan \theta_p \\ & O_{n-2p,p} & \end{pmatrix}, \right. \\ \left. \text{with } 0 \leq \theta_i \leq \pi \right\}; \quad (1.0.9)$$

also⁴,

$$T_{Id} Gr(p, \mathbb{F}^n) = \{Z \mid \text{arbitrary } (n-p) \times p \text{ matrix, with } Z_{ij} \in \mathbb{F}\}. \quad (1.0.10)$$

⁴Setting $\text{Id} = \begin{pmatrix} I_p \\ O \end{pmatrix}$.

Setting, for the respective cases of $Gr(p, \mathbb{F}^n)$ and $T_{Id}Gr(p, \mathbb{F}^n)$,

$$\begin{aligned} \{y_1, \dots, y_p\} &= \text{spectrum} \left(\frac{1 - Z^\dagger Z}{1 + Z^\dagger Z} \right) = (\cos 2\theta_1, \dots, \cos 2\theta_p) \\ \{u_1, \dots, u_p\} &:= \text{spectrum} (Z^\dagger Z) \end{aligned}$$

with $-1 \leq y_i \leq 1$, and $0 \leq u_i < \infty$, Haar measure on $Gr(p, \mathbb{F}^n)$ and $T_{Id}Gr(p, \mathbb{F}^n)$ reads, setting $k = n - 2p$, (**Weyl integration formulae**)

$$\begin{aligned} Gr(p, \mathbb{F}^n) : \quad d\mu(Z) &= |\Delta_p(y)|^{2\beta} \prod_{i=1}^p (1 - y_i)^{\beta k + (\beta - 1)} (1 + y_i)^{\beta - 1} dy_i dK \\ T_{Id}Gr(p, \mathbb{F}^n) : \quad d\mu(Z) &= |\Delta_p(u)|^{2\beta} \prod_{i=1}^p u_i^{\beta k + (\beta - 1)} du_i dK, \end{aligned} \quad (1.0.11)$$

leading to the table:

G/K	induced measure $d\mu$ on $K \backslash G/K$	$d\mu$ on $T_{Id}G/K$
$Gr(p, \mathbb{C}^n)$	$\Delta_p(y)^2 \prod_1^p (1 - y_i)^k dy_i$	$\Delta_p(u)^2 \prod_1^p u_i^k du_i$
$Gr(p, \mathbb{R}^n)$	$ \Delta_p(y) \prod_1^p (1 - y_i)^{\frac{1}{2}(k-1)} (1 + y_i)^{-\frac{1}{2}} dy_i$	$ \Delta_p(u) \prod_1^p u_i^{\frac{1}{2}(k-1)} du_i$
$Gr(p, \mathbb{H}^n)$	$\Delta_p(y)^4 \prod_1^p (1 - y_i)^{2k+1} (1 + y_i) dy_i$	$\Delta_p(u)^4 \prod_1^p u_i^{2k+1} du_i$

Table 1

In the other description (1.0.4) of $Gr(p, \mathbb{F}^n)$, given by

$$Gr(p, \mathbb{F}^n) \simeq \mathcal{S} := \left\{ M = AI_{p, n-p} A^{-1} I_{p, n-p} \mid A \in G \right\}$$

an appropriate left action of $B \in K$ on $A \in G$, amounting to conjugation in $\mathcal{S}$, leads to the matrix in the torus $\mathfrak{A} \subset G$,

$$\begin{aligned} (BA)I_{p, n-p}(BA)^{-1}I_{p, n-p} &= B(AI_{p, n-p}A^{-1}I_{p, n-p})B^{-1} \\ &= \begin{pmatrix} \Re e^{2i\Theta_p} & \Im e^{2i\Theta_p} & & \\ -\Im e^{2i\Theta_p} & \Re e^{2i\Theta_p} & & \\ & & O & \\ & & & I_{n-2p} \end{pmatrix} \in \mathfrak{A}, \end{aligned}$$

with $\Theta_p := \text{diag}(\theta_1, \dots, \theta_p)$.

Proof of Proposition 1.4: The p columns of the $n \times p$ matrix $A \begin{pmatrix} I_p \\ O \end{pmatrix}$, with $A \in G$, and $O := O_{n-p,p}$ (a zero matrix of size $(n-p, p)$) span a p -dimensional plane in $\mathbb{F}^n$, so that

$$Gr(p, \mathbb{F}^n) = \left\{ \text{span } A \begin{pmatrix} I_p \\ O \end{pmatrix}, \text{ with } A \in G \right\}.$$

The **right action** of $B \in K$ on $A \in G$ acts on the p -plane $A \begin{pmatrix} I_p \\ O \end{pmatrix}$ as:

$$AB \begin{pmatrix} I_p \\ O \end{pmatrix} = A \begin{pmatrix} B_{11} & O \\ O & B_{22} \end{pmatrix} \begin{pmatrix} I_p \\ O \end{pmatrix} = A \begin{pmatrix} B_{11} \\ O \end{pmatrix} = A \begin{pmatrix} I_p \\ O \end{pmatrix} B_{11}$$

and therefore it has no effect on that plane

$$\text{span } AB \begin{pmatrix} I_p \\ O \end{pmatrix} = \text{span } A \begin{pmatrix} I_p \\ O \end{pmatrix} B_{11} = \text{span } A \begin{pmatrix} I_p \\ O \end{pmatrix},$$

since multiplication to the right by B_{11} merely replaces the p columns of $A \begin{pmatrix} I_p \\ O \end{pmatrix}$ by p linear combination. Then the $n \times p$ matrix $V := A \begin{pmatrix} I_p \\ O \end{pmatrix}$ satisfies

$$V^\dagger V = (I_p \ O) A^\dagger A \begin{pmatrix} I_p \\ O \end{pmatrix} = (I_p \ O) I_n \begin{pmatrix} I_p \\ O \end{pmatrix} = I_p. \quad (1.0.12)$$

Conversely, we show that

$$\text{span } A_1 \begin{pmatrix} I_p \\ O \end{pmatrix} = \text{span } A_2 \begin{pmatrix} I_p \\ O \end{pmatrix} \quad (1.0.13)$$

implies $A_2^{-1} A_1 \in K = K_1 \times K_2$. Indeed, (1.0.13) holds if and only if

$$A_1 \begin{pmatrix} I_p \\ O \end{pmatrix} = A_2 \begin{pmatrix} I_p \\ O \end{pmatrix} g, \text{ with an invertible } p \times p \text{ matrix } g. \quad (1.0.14)$$

Then we prove $g \in K_1$. Indeed, from (1.0.13), the matrices $V_i := A_i \begin{pmatrix} I_p \\ O \end{pmatrix}$ satisfy $V^\dagger V = I_p$ and so $V_1 = V_2 g$ implies $g^\dagger g = (V_2 g)^\dagger V_2 g = V_1^\dagger V_1 = I_p$. Multiplying (1.0.14) to the left with A_2^{-1} yields

$$A_2^{-1} A_1 \begin{pmatrix} I_p \\ O \end{pmatrix} = \begin{pmatrix} g \\ O \end{pmatrix} \quad \text{and thus } G \ni A_2^{-1} A_1 = \begin{pmatrix} g & * \\ O & h \end{pmatrix};$$

the fact that the latter matrix is in G , implies $* = 0$ and $g \in K_1$, $h \in K_2$. This means that $A_2^{-1} A_1 \in K$ and so $Gr(p, \mathbb{F}^n) \cong G/K$.

To describe Affine $Gr(p, \mathbb{F}^n)$, notice that a plane $A \begin{pmatrix} I_p \\ O \end{pmatrix}$ for which $\det A_{11} \neq 0$, can be expressed as

$$G/K \ni \text{span } A \begin{pmatrix} I_p \\ O \end{pmatrix} = \text{span} \begin{pmatrix} A_{11} \\ A_{21} \end{pmatrix} = \text{span} \begin{pmatrix} A_{11} \\ A_{21} \end{pmatrix} A_{11}^{-1} = \text{span} \begin{pmatrix} I_p \\ Z(A) \end{pmatrix},$$

where $Z(A) := A_{21} A_{11}^{-1}$ is a $(n-p) \times p$ matrix. Also notice that $Z(A)$ is unchanged upon multiplying A to the right with $B \in K$.

The **left action** of $B \in K$ on $A \in G$ has the following effect on

$$\text{Affine } G/K \longrightarrow \text{Affine } G/K : \begin{pmatrix} I_p \\ Z \end{pmatrix} \mapsto \begin{pmatrix} I_p \\ B_{22} Z B_{11}^{-1} \end{pmatrix}, \quad (1.0.15)$$

because in

$$BA \begin{pmatrix} I_p \\ O \end{pmatrix} = \begin{pmatrix} B_{11} & O \\ O & B_{22} \end{pmatrix} \begin{pmatrix} A_{11} \\ A_{21} \end{pmatrix} = \begin{pmatrix} B_{11} A_{11} \\ B_{22} A_{21} \end{pmatrix},$$

we have $Z(BA) = (B_{22} \ A_{21})(B_{11} \ A_{11})^{-1} = B_{22} Z(A) B_{11}^{-1}$. Picking arbitrary matrices $B_{11} \in K_1$, $B_{22} \in K_2$, the $(n-p) \times p$ ($n \geq 2p$) matrix $Z(A)$ can be “diagonalized”, namely

$$Z(BA) = B_{22} Z(A) B_{11}^{-1} = \begin{pmatrix} \alpha_1 & & O \\ & \ddots & \\ O & & \alpha_p \\ & & & O \end{pmatrix}. \quad (1.0.16)$$

Now we use the fact that the $p \times p$ matrix $Z^\dagger Z$ is “self-adjoint” and positive definite⁵, and so, setting $\alpha_i =: -\tan \theta_i$,

$$(Z(BA))^\dagger Z(BA) = \begin{pmatrix} \alpha_1^2 & & O \\ & \ddots & \\ O & & \alpha_p^2 \end{pmatrix} = \begin{pmatrix} \tan^2 \theta_1 & & O \\ & \ddots & \\ O & & \tan^2 \theta_p \end{pmatrix}. \quad (1.0.17)$$

Therefore, by the left action of K on G , the p -plane in $\mathbb{F}^n$ can be represented by the span of the columns of the following matrix, which by taking the linear combination of the columns, each multiplied with $\cos \theta_i$ reads

$$\text{span} \left(\begin{array}{ccc|ccc} & & I_p & & & \\ \hline -\tan \theta_1 & & & & O & \\ & \ddots & & & & \\ O & & & -\tan \theta_p & & \\ \hline & & O_{n-2p,p} & & & \end{array} \right) = \text{span} \left(\begin{array}{ccc|ccc} \cos \theta_1 & & & & O & \\ & \ddots & & & & \\ \hline O & & \cos \theta_p & & & \\ -\sin \theta_1 & & & O & & \\ & \ddots & & & & \\ \hline O & & & -\sin \theta_p & & \\ & & O_{n-2p,p} & & & \end{array} \right),$$

and so

$$K \backslash G / K \simeq \left\{ \text{span} \left(\begin{array}{c} \Re e^{i\Theta_p} \\ -\Im e^{i\Theta_p} \\ O_{n-2p,p} \end{array} \right), 0 \leq \theta_i \leq \pi \right\}.$$

Similarly, $Z \in T_{Id}Gr(p, \mathbb{F}^n)$ can be diagonalized by the action (1.0.16) on $T_{Id}Gr(p, \mathbb{F}^n)$; i.e.,

$$Z \mapsto B_{22} Z B_{11}^{-1} = \begin{pmatrix} z_1 & & O \\ & \ddots & \\ O & & z_p \\ \hline & & O \end{pmatrix} \in \mathfrak{a}^+, \quad (1.0.18)$$

where the z_i are linearized versions of the $\tan \theta_i$'s and where $\mathfrak{a}^+$ is a fixed Weyl chamber in the Cartan of $\mathfrak{p}$.

⁵since $v^\dagger Z^\dagger Z v = (Zv)^\dagger Z v = \sum_1^p |(Zv)_i|^2 > 0$, for $v \in \mathbb{F}^p \setminus 0$.

By the Weyl integration formula, the measure induced on $K \backslash G / K$, via the Haar measure and the embedding $G / K \hookrightarrow G$, is given by (in the compact case) (see Helgason [22], p. 188)

$$d\mu(H) = \prod_{\alpha \in \Sigma^+} |\sin \alpha(iH)|^{m_\alpha} dH, \quad H \in \mathfrak{a}_*,$$

where $g = \mathfrak{k} + \mathfrak{p}$, with compact real form $u = \mathfrak{k} + \mathfrak{p}_*$, $\mathfrak{p}_* = i\mathfrak{p}$ and with $\mathfrak{a}_*$ a maximal abelian subspace of $\mathfrak{p}$, where Σ^+ is the set of roots having positive values on the fixed Weyl chamber $\mathfrak{a}^+$ of $\mathfrak{a}_*$ and where the root space g_α has dimension m_α for any restricted root α :

$$g_\alpha := \{X \in g \mid [H, X] = \alpha(H)X, \text{ for all } H \in \mathfrak{a}\},$$

and we also have the induced measure on $T_{Id}(K \backslash G / K)$ ([22], p. 195)

$$d\mu = \prod_{\alpha \in \Sigma^+} \alpha(H)^{m_\alpha} dH, \quad H \in \mathfrak{a}^+.$$

The roots and multiplicities m_α are as follows ($n \geq 2p$):

$Gr(p, \mathbb{C}^n)$	$Gr(p, \mathbb{R}^n)$	$Gr(p, \mathbb{H}^n)$	$\alpha \in \Sigma^+$
2	1	4	$i(\varepsilon_j + \varepsilon_k)$
2	1	4	$i(\varepsilon_j - \varepsilon_k)$
1	0	3	$2i\varepsilon_\ell$
$2(n - 2p)$	$n - 2p$	$4(n - 2p)$	$i\varepsilon_\ell$

with $1 \leq j < k \leq p$, $1 \leq \ell \leq p$, yielding Table 1, upon setting

$$H = i(\theta_1, \dots, \theta_p), 0 \leq \theta_i \leq \pi, \quad (1.0.19)$$

e.g., we check table 1 for $Gr(p, \mathbb{H}^n)$. Setting $k = n - 2p$ and $y_j = \cos 2\theta_j$, which is very natural in view of (1.0.20):

for $K \backslash G / K$:

$$\begin{aligned}
d\mu &= \prod_{\alpha \in \Sigma^+} |\sin \alpha(iH)|^{m_\alpha} dH \\
&= \prod_{1 \leq j < k \leq p} |\sin(\theta_j - \theta_k) \sin(\theta_j + \theta_k)|^4 \prod_{j=1}^p |\sin 2\theta_j|^3 |\sin^2 \theta_j|^{2(n-2p)} d\theta_j \\
&= \prod_{1 \leq j < k \leq p} \left| \frac{1}{2} (\cos 2\theta_j - \cos 2\theta_k) \right|^4 \prod_{j=1}^p (1 - \cos^2 2\theta_j) \left(\frac{1 - \cos 2\theta_j}{2} \right)^{2k} \frac{1}{2} d \cos 2\theta_j \\
&= 2^{-p(2(p+k)-1)} \Delta(y)^4 \prod_{j=1}^p (1 + y_j)(1 - y_j)^{2k+1} dy_j
\end{aligned} \tag{1.0.20}$$

for $T_{id}K \backslash G / K$:

$$\begin{aligned}
d\mu &= \prod_{\alpha \in \Sigma^+} \alpha(H)^{m_\alpha} dH \\
&= c \prod_{1 \leq j < k \leq p} (v_j - v_k)^4 (v_j + v_k)^4 \prod_{j=1}^p v_j^{4k+3} dv_j \\
&= c \prod_{1 \leq j < k \leq p} (v_j^2 - v_k^2)^4 \prod_{j=1}^p (v_j^2)^{2k+1} \frac{1}{2} dv_j^2 \\
&= c 2^{-p} \Delta^4(u) \prod_{j=1}^p u_j^{2k+1} du_j,
\end{aligned} \tag{1.0.21}$$

setting $u_j = v_j^2$, where in the above we made the identification (1.0.6) of Proposition 1.3 and put Z in the normal form (1.0.16), so that $Z^\dagger Z = \text{diag}(u_1, \dots, u_p)$.

To describe G/K in a second way, remember $g = \mathfrak{k} + \mathfrak{p}$, with $\mathfrak{k}, \mathfrak{p}$ the $\pm$ eigenspaces of a lie algebra involution σ . The latter lifts to the group as an involution σ , which commutes with inversion, i.e., $(g^\sigma)^{-1} = (g^{-1})^\sigma$. Use σ to define the following embedding

$$\iota : G \hookrightarrow G : g \mapsto \iota(g) = g(g^\sigma)^{-1}, \tag{1.0.22}$$

which induces a natural injective map

$$\iota : G/K \hookrightarrow G : g \mapsto \iota(g).$$

Indeed, $\iota(g_1) = \iota(g_2)$ is equivalent to $g_1(g_1^\sigma)^{-1} = g_2(g_2^\sigma)^{-1}$, which amounts to $g_2^{-1}g_1 = (g_2^\sigma)^{-1}g_1^\sigma = (g_2^{-1})^\sigma g_1^\sigma = (g_2^{-1}g_1)^\sigma$, meaning that $g_2^{-1}g_1 \in K$.

From the polar decomposition $G = K\mathcal{A}K$, we have that every $g \in G$ can be decomposed into

$$g = k_1^{-1}A(\theta)k_2^{-1}, \quad k_1, k_2 \in K, \quad A(\theta) \in \mathcal{A},$$

with θ the torus coordinates, such that $A(\theta)A(\theta') = A(\theta + \theta')$, $A(0) = I$ and $A^{-1}(\theta) = A(-\theta)$. Since $A = \exp a$, with $a \in i\mathfrak{p}$, we have $A(\theta)^\sigma = A^{-1}(\theta)$. From (1.0.22) the torus $\mathcal{A}$ embeds into $\mathcal{S}$ as follows:

$$\iota(A(\theta)) = A(\theta)(A^\sigma(\theta))^{-1} = A(\theta)A(\theta) = A(2\theta).$$

Moreover the polar decomposition $g = k_1^{-1}A(\theta)k_2^{-1}$ yields conjugation in $\mathcal{S}$ by k_1 :

$$\begin{aligned} \iota(g) &= k_1^{-1}(k_1 g k_2 k_2^{-1} (g^\sigma)^{-1} k_1^{-1}) k_1 \\ &= k_1^{-1}(k_1 g k_2 (k_2^\sigma)^{-1} (g^\sigma)^{-1} (k_1^\sigma)^{-1}) k_1 \\ &= k_1^{-1}(k_1 g k_2 ((k_1 g k_2)^\sigma)^{-1}) k_1 \\ &= k_1^{-1} \iota(k_1 g k_2) k_1 \\ &= k_1^{-1} \iota(A(\theta)) k_1 \\ &= k_1^{-1} A(2\theta) k_1 \end{aligned} \tag{1.0.23}$$

Specializing to the Grassmannian case, we have the involution

$$\sigma : G \rightarrow G : g \mapsto g^\sigma := I_{p,n-p} g I_{p,n-p}$$

with

$$\begin{aligned} K &= \{\text{fixed points in } G \text{ of the involution } \sigma\} \\ &= \{g \in G \text{ such that } g I_{p,n-p} = I_{p,n-p} g\}. \end{aligned}$$

Setting

$$\Theta_p := \text{diag}(\theta_1, \dots, \theta_p),$$

we have that a maximal abelian subspace of the Lie algebra is given by

$$\mathfrak{a} = \left\{ a(\theta) = \begin{pmatrix} O_p & \Theta_p & O \\ -\Theta_p & O_p & O \\ O & O & I_{n-2p} \end{pmatrix}, \quad 0 \leq \theta < 2\pi \right\}$$

and by exponentiation, we find the torus

$$\mathfrak{A} = \left\{ A(\theta) = \begin{pmatrix} \Re e^{i\theta_p} & \Im e^{i\theta_p} & & \\ -\Im e^{i\theta_p} & \Re e^{i\theta_p} & O & \\ & O & & I_{n-2p} \end{pmatrix}, \quad 0 \leq \theta < 2\pi \right\} \in G.$$

The spectrum of $A(\theta)$ is easily seen to be

$$(e^{i\theta_1}, \dots, e^{i\theta_p}, e^{-i\theta_1}, \dots, e^{-i\theta_p}, \underbrace{1, \dots, 1}_{n-2p}). \quad (1.0.24)$$

To connect with the previous description, given $g \in G$, we pick k_1, k_2 such that $k_1 g k_2 = A(\theta)$ and by the previous discussions, we have

$$\begin{aligned} Z^\dagger(g)Z(g) &= Z^\dagger(k_1 g k_2)Z(k_1 g k_2) \\ &= Z^\dagger(A(\theta))Z(A(\theta)) \\ &= \text{diag}(\tan^2 \theta_1, \dots, \tan^2 \theta_p). \end{aligned}$$

The embedding $\mathfrak{A} \hookrightarrow Gr(p, \mathbb{F}^n)$ is then given by

$$\mathfrak{A} \hookrightarrow Gr(p, \mathbb{F}^n) : A(\theta) \mapsto A(\theta) \begin{pmatrix} I_p \\ O_{n-p,p} \end{pmatrix} = \begin{pmatrix} \Re e^{i\theta_p} \\ -\Im e^{i\theta_p} \\ O_{n-2p,p} \end{pmatrix},$$

and so the θ_i 's in the two discussions are identical. ■

Proof of Theorems 1.1, 1.2 and 1.3: In order to compute integral (1.0.3), recall in the above description of $K \backslash G / K$,

$$H = i(\theta_1, \dots, \theta_p).$$

Since the integrand is invariant under the left action of K on G , which induces on Z the map $Z \rightarrow B_{22} Z B_{11}^{-1}$, use (1.0.17), from which it follows that, upon using $\tan^2 \theta_i = \frac{1 - \cos 2\theta_i}{1 + \cos 2\theta_i} = \frac{1 - y_i}{1 + y_i}$, and $\cos^2 \theta_i = (1 + y_i)/2$, we have

$$Z^\dagger Z = \text{diag}(\tan^2 \theta_1, \dots, \tan^2 \theta_p) = \text{diag}\left(\frac{1 - y_1}{1 + y_1}, \dots, \frac{1 - y_p}{1 + y_p}\right)$$

and

$$(I + Z^\dagger Z)^{-1} = \begin{pmatrix} \cos^2 \theta_1 & & O \\ & \ddots & \\ O & & \cos^2 \theta_p \end{pmatrix} = \frac{1}{2} \operatorname{diag}(1 + y_1, \dots, 1 + y_p).$$

Hence, setting $z_i = \cos^2 \theta_i$ in the last identity below, we have $y_i = 2z_i - 1$, using (1.0.11) and picking an appropriate normalizing constant c :

$$\begin{aligned} & \int_{Gr(p, \mathbb{F}^n)} e^{x \operatorname{Tr}(I + Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-\beta(q-p)} d\mu(Z) \\ &= c \int_{[-1, 1]^p} e^{\frac{x}{2} \sum_1^p (1 + y_i)} |\Delta_p(y)|^{2\beta} \prod_{i=1}^p \left(\frac{1 - y_i}{1 + y_i} \right)^{\beta(p-q)} (1 - y_i)^{\beta(n-2p)} (1 - y_i^2)^{\beta-1} dy_i \\ &= c \int_{[0, 1]^p} e^{x \sum_1^p z_i} |\Delta_p(z)|^{2\beta} \prod_1^p (1 - z_i)^{\beta(n-p-q+1)-1} z_i^{\beta(q-p+1)-1} dz_i, \end{aligned}$$

upon setting $y_i = 2z_i - 1$.

Finally, using the Weyl integration formula (1.0.21) on the tangent space and identifying the $(z_1, \dots, z_p)$ in (1.0.18) with a point in $\mathfrak{a}^+$, and setting $u_i = z_i^2 \geq 0$, $1 \leq i \leq p$, we find

$$\begin{aligned} & \int_{\substack{Z \in T_{id} Gr(p, \mathbb{F}^n) \\ \text{with spectrum } (Z^\dagger Z) \in E}} e^{-x \operatorname{Tr} Z^\dagger Z} \det(Z^\dagger Z)^{\beta p} d\mu(Z), \\ &= \int_{E^p} |\Delta_p(u)|^{2\beta} \prod_1^p e^{-x u_i} u_i^{\beta(n-p+1)-1} du_i, \end{aligned}$$

establishing Theorem 1.3.

Finally, to prove Theorem 1.2, notice that, according to (1.0.23), matrices M in $\mathcal{S}$ can be diagonalized to matrices $A(2\theta)$, with spectrum

$$(e^{2i\theta_1}, \dots, e^{2i\theta_p}, e^{-2i\theta_1}, \dots, e^{-2i\theta_p}, \underbrace{1, \dots, 1}_{n-2p}). \quad (1.0.25)$$

So M decomposes into $M = M_0 \otimes M_1$, with M_1 being the $n - 2p$ -dimensional eigenspace corresponding to the eigenvalue 1 and so

$$\prod_1^p \frac{1 - y_k}{1 + y_k} = \prod_1^p \frac{1 - \cos 2\theta_k}{1 + \cos 2\theta_k} = \prod_1^p \frac{(1 - e^{2i\theta_k})(1 - e^{-2i\theta_k})}{(1 + e^{2i\theta_k})(1 + e^{-2i\theta_k})} = \det \frac{I - M_0}{I + M_0}.$$

■

2 Jack polynomials

2.1 Young diagrams and Schur polynomials

Standard references to this subject are MacDonald, Sagan, Stanley, Stanton and White [34, 41, 44, 45]. To set the notation, we remind the reader of a few basic facts.

- A *partition* of $n = |\lambda| := \lambda_1 + \dots + \lambda_\ell$ (with $n = |\lambda|$ called the weight) is represented by a *Young diagram* $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_\ell \geq 0$. A *dual Young diagram* $\lambda^\top = (\lambda_1^\top \geq \lambda_2^\top \geq \dots)$ is the diagram obtained by flipping the diagram λ about its diagonal.
- A *semi-standard Young tableau* of shape λ is an array of positive integers a_{ij} placed at (i, j) in the Young diagram λ , which are non-decreasing from left to right *and* strictly increasing from top to bottom.
- A *standard Young tableau* of shape λ is an array of integers $1, \dots, n$ placed in the Young diagram, which are strictly increasing from left to right *and* from top to bottom.
- The *Schur polynomial* s_λ associated with a Young diagram λ is a symmetric function in the variables $x_1, x_2, \dots$, (finite or infinite), where $n = |\lambda|$ and defined by (for notation f^λ , see the next point)

$$s_\lambda(x_1, x_2, \dots) = \sum_{\substack{\{a_{i,j}\} \text{ semi-standard} \\ \text{tableaux } \lambda}} \prod_{(i,j) \in \lambda} x_{a_{i,j}} = f^\lambda x_1 \dots x_n + \dots$$

- The *hook length* of the i, j th box is defined by $h_{ij}^\lambda := \lambda_i + \lambda_j^\top - i - j + 1$. Also define

$$\begin{aligned} h^\lambda &:= \prod_{(i,j) \in \lambda} h_{ij}^\lambda \\ &= \frac{\prod_1^m (m + \lambda_i - i)!}{\Delta_m(m + \lambda_1 - 1, \dots, m + \lambda_m - m)}, \quad \text{for } m \geq \lambda_1^\top \end{aligned} \quad (2.1.1)$$

- The *number of standard Young tableaux* of a given shape $\lambda = (\lambda_1 \geq$

$\dots \geq \lambda_m)$ is given by

$$\begin{aligned}
f^\lambda &= \#\{\text{standard tableaux of shape } \lambda\} \\
&= |\lambda|! \left| s_\lambda(x) \right|_{\substack{\sum_k x_k = 1 \\ \sum_k x_k^i = 0 \text{ for } i \geq 2}} \\
&= \text{coefficient of } x_1 x_2 \dots x_n \text{ in } s_\lambda(x) \\
&= \frac{|\lambda|!}{h^\lambda} = |\lambda|! \det \left(\frac{1}{(\lambda_i - i + j)!} \right) \\
&= |\lambda|! \frac{\Delta_m(m + \lambda_1 - 1, \dots, m + \lambda_m - m)}{\prod_1^m (m + \lambda_i - i)!}, \quad \text{for } m \geq \lambda_1^\top.
\end{aligned} \tag{2.1.2}$$

- The *number of semi-standard Young tableaux* of a given shape λ , with numbers 1 to k for $k \geq 1$:

$$\begin{aligned}
&\# \left\{ \begin{array}{l} \text{semi-standard tableaux of shape } \lambda \\ \text{filled with numbers from 1 to } k \end{array} \right\} \\
&= s_\lambda(\overbrace{1, \dots, 1}^k, 0, 0, \dots) \\
&= \prod_{(i,j) \in \lambda} \frac{j - i + k}{h_{i,j}^\lambda} \\
&= \begin{cases} \frac{\Delta_k(k + \lambda_1 - 1, \dots, k + \lambda_k - k)}{\prod_{i=1}^{k-1} i!}, & \text{when } k \geq \lambda_1^\top, \\ 0, & \text{when } k < \lambda_1^\top, \end{cases}
\end{aligned} \tag{2.1.3}$$

using the fact that

$$\prod_{(i,j) \in \lambda} (j - i + k) = \frac{\prod_{i=1}^k (k + \lambda_i - i)!}{\prod_1^{k-1} i!}. \tag{2.1.4}$$

- *Robinson-Schensted-Knuth correspondence*: Given

$$\begin{aligned} S_n &= \text{group of permutations of } \{1, \dots, n\} \\ S_n^k &= \{\text{words of length } n \text{ built from the set } \{1, \dots, k\}\} \end{aligned}$$

the following 1-1 correspondences hold:

$$S_n \longrightarrow \left\{ \begin{array}{l} (P, Q), \text{ where } P \text{ and } Q \text{ are two standard Young tableaux} \\ \text{of same shape } \lambda, \text{ with } |\lambda| = n \text{ and taken from } \{1, \dots, n\} \end{array} \right\}$$

$$S_n^k \longrightarrow \left\{ \begin{array}{l} (P, Q), \text{ where } P \text{ and } Q \text{ have same shape } \lambda, \text{ with } |\lambda| = n \\ P \text{ is semi-standard, filled with numbers from } 1 \text{ to } k, \text{ and} \\ Q \text{ is standard, filled with numbers from } 1 \text{ to } n \end{array} \right\}$$

It follows that for given n and k , we have

$$\sum_{\substack{\lambda \text{ with} \\ |\lambda| = n}} (f^\lambda)^2 = n! \quad (2.1.5)$$

$$\sum_{\substack{\lambda \text{ with} \\ |\lambda| = n}} f^\lambda s_\lambda(1^k) = k^n \quad (2.1.6)$$


- *Increasing and decreasing sequences*

According to Greene [20], given a word $\pi \in S_n^k$, mapped, via the RSK correspondence, into (P, Q) of shape $\lambda = (\lambda_1, \dots, \lambda_\ell)$, then for any k ,

$$\begin{aligned} \lambda_1 + \dots + \lambda_k &= \left\{ \begin{array}{l} \text{length of the longest } \textit{weakly} \\ k\text{-increasing subsequence} \end{array} \right\} \\ \lambda_1^\top + \dots + \lambda_k^\top &= \left\{ \begin{array}{l} \text{length of the longest } \textit{strictly} \\ k\text{-decreasing subsequence} \end{array} \right\} \end{aligned}$$

2.2 Some useful formulae on hook length

Remembering the notation (0.0.8), we have the following statement:



Lemma 2.1 *Given a partition $\lambda \supseteq \mu := (n-p)^p$ with $\lambda_1^\top = p$, $0 \leq p < n$, then⁶*

$$h^\lambda = h^\mu h^{\lambda \setminus \mu} \frac{(n)_{\lambda \setminus \mu}}{(p)_{\lambda \setminus \mu}}, \quad (2.2.1)$$

$$h^\mu = \frac{\prod_{i=1}^p (n-i)!}{p-1} \quad \text{and} \quad s_\mu(1^p) = 1, \quad (2.2.2)$$

$$\frac{s_\lambda(1^p)}{h^\lambda} = \frac{1}{h^\mu (h^{\lambda \setminus \mu})^2} \frac{((p)_{\lambda \setminus \mu})^2}{(n)_{\lambda \setminus \mu}}. \quad (2.2.3)$$

Proof: Setting $\kappa = \lambda \setminus \mu$

$$\begin{aligned} \frac{h^\lambda}{h^{\lambda \setminus \mu}} &= \prod_{(i,j) \in \mu} h_{(i,j)}^\lambda \\ &= \prod_{i=1}^p \prod_{j=1}^{n-p} (n+1+\kappa_i-i-j) \\ &= \prod_{i=1}^p (n+\kappa_i-i) \dots (n+1+\kappa_i-i-n+p) \\ &= h^\mu \prod_{i=1}^p \frac{(p+1+\kappa_i-i) \dots (n+\kappa_i-i)}{(p+1-i) \dots (n-i)} \\ &= h^\mu \prod_{i=1}^p \frac{(n-i+1)_{\kappa_i}}{(p-i+1)_{\kappa_i}} \end{aligned}$$

⁶Here $(a)_\lambda := (a)_\lambda^{(1)} = \prod_i (a+1-i)_{\lambda_i}$.

$$\begin{aligned}
h^\mu &= [(n-1)(n-2)\dots p][(n-2)\dots(p-1)]\dots[(n-p)\dots 1] \\
&= \frac{(n-1)!(n-2)!\dots(n-p)!}{(p-1)!(p-2)!\dots 1!} \\
&= \frac{\prod_1^p (n-i)!}{1^{p-1} \prod_1^{p-1} i!}. \tag{2.2.4}
\end{aligned}$$

Then

$$s_\mu(1^p) = \frac{\prod_{(i,j) \in \mu} (j-i+p)}{h^\mu} = \frac{\prod_1^p (n-i)!}{h^\mu \prod_1^{p-1} i!} = 1, \quad \text{using (2.1.4).}$$

Finally, using in the second identity below formula (2.1.4) and in the fourth identity $\prod_1^p (n-i+\kappa_i)! = \prod_1^p (n-i)! \prod_1^p (n-i+1)_{\kappa_i}$, one computes

$$\begin{aligned}
\frac{s_\lambda(1^p)}{h^\lambda} &= \frac{\prod_{(i,j) \in \lambda} (j-i+p)}{(h^\lambda)^2} \\
&= \frac{1}{(h^\lambda)^2} \frac{\prod_{i=1}^p (p+\lambda_i-i)!}{\prod_1^{p-1} i!} \\
&= \frac{1}{(h^\lambda)^2} \frac{\prod_{i=1}^p (n-i+\kappa_i)!}{\prod_1^{p-1} i!}, \quad \text{since } \lambda_i = n-p+\kappa_i
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(h^\lambda)^2} \frac{\prod_{i=1}^p (n-i)!}{\prod_{i=1}^{p-1} i!} \prod_{i=1}^p (n-i+1)_{\kappa_i} \\
&= \frac{h^\mu}{(h^\mu h^{\lambda \setminus \mu})^2} \prod_{i=1}^p \frac{((p-i+1)_{(\lambda \setminus \mu)_i})^2}{(n-i+1)_{(\lambda \setminus \mu)_i}}, \quad \text{using (2.2.1) and (2.2.2)}
\end{aligned}$$

ending the proof of Lemma 2.1. ■

2.3 Jack polynomials

Define symmetric polynomials

$$\begin{aligned}
p_\lambda(x_1, x_2, \dots) &:= p_{\lambda_1} p_{\lambda_2} \cdots = \sum_i x_i^{\lambda_1} \sum_i x_i^{\lambda_2} \cdots \\
m_\lambda(x_1, x_2, \dots) &:= \sum_{\substack{\text{permutations} \\ \text{of } x_1, \dots, x_n}} x_1^{\lambda_1} \cdots x_n^{\lambda_n},
\end{aligned}$$

and the dominance ordering between partitions:

$$\mu \leq \lambda \quad \text{means} : \quad \sum_1^\ell \mu_i \leq \sum_1^\ell \lambda_i, \quad \text{for all } \ell.$$

Given that λ has $m_i = m_i(\lambda)$ parts equal to i , define the inner-product $\langle \cdot, \cdot \rangle$ on the vector space of all symmetric functions of bounded degree (to be explained)

$$\langle p_\lambda, p_\mu \rangle = \delta_{\lambda\mu} (1^{m_1} 2^{m_2} \cdots) m_1! m_2! \cdots \alpha^{\lambda^\top}.$$

Jack polynomials are the unique symmetric functions $J_\lambda^{(\alpha)}$ satisfying

- (i) $\langle J_\lambda^{(\alpha)}, J_\mu^{(\alpha)} \rangle = 0$, if $\lambda \neq \mu$,
- (ii) $J_\lambda^{(\alpha)} = \sum_{\mu \leq \lambda} v_{\lambda\mu}(\alpha) m_\mu$,
- (iii) If $|\lambda| = n$, then

$$J_\lambda^{(\alpha)} = n! x_1 \cdots x_n + \cdots$$

It follows that

$$v_{\lambda\lambda} = \prod_{(i,j) \in \lambda} (\lambda_j^\top - i + 1 + \alpha(\lambda_i - j)).$$

Special cases:

Jack polynomials for $\alpha = 1$ are proportional to Schur polynomials, namely

$$J_\lambda^{(1)} = h^\lambda s_\lambda.$$

Zonal polynomials are given by

$$Z_\lambda^{(\beta)} = J_\lambda^{(1/\beta)}, \quad \text{with } \beta = 1, 1/2, 2.$$

They have the remarkable property that for $G = O(n)$, $U(n)$ or $U(n, \mathbb{H}) := \{g \mid g\bar{g}^\top = I\}$,

$$\int_G J_\lambda^{(\alpha)}(\sigma k \tau k^{-1}) dk = \frac{J_\lambda^{(\alpha)}(\sigma) J_\lambda^{(\alpha)}(\tau)}{J_\lambda^{(\alpha)}(1^n)}, \quad (2.3.1)$$

for all

$$\begin{aligned} \sigma, \tau \in \Sigma &= \{\text{real symmetric matrices}\} && \text{for } \alpha = 2 \\ &= \{\text{Hermitian matrices}\} && \text{for } \alpha = 1 \\ &= \{\text{quaternionic matrices, with } \sigma = \bar{\sigma}^\top\} && \text{for } \alpha = 1/2. \end{aligned}$$

The function $J_\lambda^{(\alpha)}(\tau)$ is a symmetric function of the (real) spectrum of τ .

Orthogonality:

$$\langle J_\lambda^{(\alpha)}, J_\mu^{(\alpha)} \rangle = \delta_{\mu\lambda} j_\lambda^{(\alpha)}, \quad (2.3.2)$$

where

$$\begin{aligned} j_\lambda^{(\alpha)} &= \prod_{(i,j) \in \lambda} (\lambda_j^\top - i + \alpha(\lambda_i - j + 1)) (\lambda_j^\top - i + 1 + \alpha(\lambda_i - j)) \\ &= \begin{cases} (h^\lambda)^2 & \text{for } \alpha = 1 \\ h^{2\lambda} & \text{for } \alpha = 2 \\ h^{2\lambda^\top} / 2^{2|\lambda|} & \text{for } \alpha = 1/2. \end{cases} \end{aligned} \quad (2.3.3)$$

Special values:

For arbitrary n , we have

$$\begin{aligned}
J_\lambda^{(\alpha)}(1^n) &= \prod_{(i,j) \in \lambda} (n - (i-1) + \alpha(j-1)), \text{ where } 1^n = (\overbrace{1, \dots, 1}^n, 0, 0, \dots) \\
&= \alpha^{|\lambda|} \prod_{(i,j) \in \lambda} \left(\frac{1}{\alpha} (n - i + 1) + j - 1 \right) = \alpha^{|\lambda|} \left(\frac{n}{\alpha} \right)_\lambda^{(\alpha)} \\
&= \begin{cases} \alpha^{|\lambda|} \prod_{i=1}^m \frac{\Gamma\left(\frac{1}{\alpha}(n - i + 1) + \lambda_i\right)}{\Gamma\left(\frac{1}{\alpha}(n - i + 1)\right)} > 0, & \text{for all } m \geq \lambda_1^\top, \text{ if } n \geq \lambda_1^\top, \\ 0, & \text{if } n < \lambda_1^\top \end{cases} \quad (2.3.4)
\end{aligned}$$

and so, for $\alpha = 1$,

$$\begin{aligned}
s_\lambda(1^n) &= \frac{1}{h^\lambda} J_\lambda^1(1^n) \\
&= \begin{cases} \frac{1}{h^\lambda} \prod_{i=1}^m \frac{(n - i + \lambda_i)!}{(n - i)!} \geq 0, & \text{for all } m \geq \lambda_1^\top, \text{ if } n \geq \lambda_1^\top, \\ 0, & \text{if } n < \lambda_1^\top. \end{cases} \quad (2.3.5)
\end{aligned}$$

The last identity in (2.3.4) is obtained by taking the product over the i th row of λ and using $(x+n)(x+n-1)\dots x = \frac{\Gamma(x+n+1)}{\Gamma(x)}$. When the Gamma functions blow up, the formulas must be understood as limits. Also

$$\begin{aligned}
J_\lambda(x)|_{\sum_\ell x_\ell^i = \delta_{1i}u} &= u^{|\lambda|} \\
s_\lambda(x)|_{\sum_\ell x_\ell^i = \delta_{1i}u} &= \frac{u^{|\lambda|}}{h^\lambda}.
\end{aligned}$$

Expansion of $(x_1 + x_2 + \dots)^n$:

$$\frac{(x_1 + x_2 + \dots)^n}{n!} = \alpha^n \sum_{|\lambda|=n} \frac{J_\lambda^{(\alpha)}(x)}{j_\lambda^{(\alpha)}}. \quad (2.3.6)$$

Then also

$$(x_1 + x_2 + \dots)^n = \sum_{|\lambda|=n} C_\lambda^{(\alpha)}(x), \text{ with } C_\lambda^{(\alpha)}(x) = \frac{|\lambda|! \alpha^{|\lambda|}}{j_\lambda^{(\alpha)}} J_\lambda^{(\alpha)}(x). \quad (2.3.7)$$

Cauchy identity:

$$\prod_{(i,j) \geq 1} (1 - x_i y_j)^{-1/\alpha} = \sum_{\lambda \in \mathbb{Y}} \frac{J_\lambda^{(\alpha)}(x) J_\lambda^{(\alpha)}(y)}{j_\lambda^{(\alpha)}}; \quad (2.3.8)$$

in particular, for $\alpha = 1$

$$\prod_{i,j \geq 1} (1 - x_i y_j)^{-1} = \sum_{\lambda \in \mathbb{Y}} s_\lambda(x) s_\lambda(y). \quad (2.3.9)$$

Hypergeometric functions

Generalized hypergeometric functions ${}_2F_1^{(\alpha)}$ are defined by:

$${}_2F_1^{(\alpha)}(p, q; n; x) := \sum_{\kappa \in \mathbb{Y}} \frac{(p)_\kappa^{(\alpha)} (q)_\kappa^{(\alpha)}}{(n)_\kappa^{(\alpha)}} \alpha^{|\kappa|} \frac{J_\kappa^{(\alpha)}(x)}{j_\kappa^{(\alpha)}}. \quad (2.3.10)$$

For $\alpha = 1$, using $J_\kappa^{(1)} = h^\kappa s_\kappa$ and $j_\kappa^{(1)} = (h^\kappa)^2$, we have

$${}_2F_1^{(1)}(p, q; n; x) := \sum_{\kappa \in \mathbb{Y}} \frac{(p)_\kappa^{(1)} (q)_\kappa^{(1)}}{(n)_\kappa^{(1)}} \frac{s_\kappa(x)}{h^\kappa}, \quad (2.3.11)$$

and so, upon restriction,

$${}_2F_1^{(1)}(p, q; n; x) \Big|_{\sum_\ell x_\ell^i = \delta_{1i} u} = \sum_{\kappa \in \mathbb{Y}} u^{|\kappa|} \frac{(p)_\kappa^{(1)} (q)_\kappa^{(1)}}{(h^\kappa)^2 (n)_\kappa^{(1)}} \quad (2.3.12)$$

Generalized Selberg formula:

Kaneko [30] computes the following integrals, subjected to the condition that $a, b > \beta(p-1)$ (see also MacDonald [34] and Kadell [29]):

$$\begin{aligned}
& \int_{[0,1]^p} J_\lambda^{(1/\beta)}(x) |\Delta_p(x)|^{2\beta} \prod_{i=1}^p (1-x_i)^{a-\beta(p-1)-1} x_i^{b-\beta(p-1)-1} dx_i \\
&= J_\lambda^{(1/\beta)}(1^p) \prod_{i=1}^p \frac{\Gamma(i\beta+1) \Gamma(a+\beta(1-i)) \Gamma(\lambda_i+b+\beta(1-i))}{\Gamma(\beta+1) \Gamma(\lambda_i+a+b+\beta(1-i))}.
\end{aligned} \tag{2.3.13}$$

Setting $u = (u_1, \dots, u_m)$, we have the following representation in terms of the hypergeometric function:

$$\begin{aligned}
& \int_{[0,1]^p} \prod_{\substack{1 \leq i \leq p \\ 1 \leq k \leq m}} (1-x_i u_k)^{-\beta} |\Delta_p(x)|^{2\beta} \prod_{i=1}^p (1-x_i)^{a-\beta(p-1)-1} x_i^{b-\beta(p-1)-1} dx_i \\
&= {}_2F_1^{(1/\beta)}(\beta p, b, a+b; u) \prod_{i=1}^p \frac{\Gamma(i\beta+1) \Gamma(a+\beta(1-i)) \Gamma(b+\beta(1-i))}{\Gamma(\beta+1) \Gamma(a+b+\beta(1-i))}.
\end{aligned} \tag{2.3.14}$$

3 Probability measures on partitions

3.1 Probability measure on the set $\mathbb{Y}$ of all partitions

In view of formula (2.3.8), define the (not necessarily positive) probability measure on the space $\mathbb{Y}$ of Young diagrams, depending on x, y and α : (see [10, 11, 12, 50, 31])

$$P(\lambda) := \frac{J_\lambda^{(\alpha)}(x) J_\lambda^{(\alpha)}(y)}{j_\lambda^{(\alpha)} \prod_{i,j} (1-x_i y_j)^{-1/\alpha}}, \quad \lambda \in \mathbb{Y}. \tag{3.1.1}$$

In particular, evaluating $P(\lambda)$ along the locus

$$\mathcal{L} = \begin{cases} \text{for } x = (x_1, x_2, \dots) \text{ such that } \sum_{\ell} x_{\ell}^i = \delta_{1i} u, \text{ we have } J_\lambda^{(\alpha)}(x) = u^{|\lambda|} \\ \text{for } y = (\overbrace{1, 1, \dots, 1}^p, 0, 0, \dots) = 1^p, \text{ we have } J_\lambda^{(\alpha)}(1^p) \text{ as in (2.3.4),} \end{cases} \tag{3.1.2}$$

and using⁷

$$\prod_{i,j \geq 1} (1 - x_i y_j)^{-1} = \prod_{j=1}^p \prod_{i \geq 1} (1 - x_i y_j)^{-1} \Big|_{y_j=1} = \prod_{i \geq 1} (1 - x_i)^{-p} = e^{pu},$$

we obtain the genuine (≥ 0) probability measure for $u > 0$ on the space $\mathbb{Y}$, depending on $u \in \mathbb{R}$ and the integer $p > 0$,

$$\begin{aligned} P_{u,p}(\lambda) &:= P(\lambda) \Big|_{\mathcal{L}} = e^{-pu/\alpha} u^{|\lambda|} \frac{J_{\lambda}^{(\alpha)}(1^p)}{j_{\lambda}^{(\alpha)}} \\ &= e^{-pu/\alpha} \frac{(pu/\alpha)^{|\lambda|}}{|\lambda|!} P^{\ell,p}(\lambda), \text{ with } \ell := |\lambda|, \end{aligned} \quad (3.1.3)$$

with

$$(\text{support } P_{u,p}) \subseteq \mathbb{Y}^{(p)} := \{\lambda \in \mathbb{Y} \text{ such that } \lambda_1^{\top} \leq p\}.$$

Probability (3.1.3) can be viewed as a Poissonized probability of

$$P^{\ell,p}(\lambda) = \frac{J_{\lambda}^{(\alpha)}(1^p) \ell!}{j_{\lambda}^{(\alpha)} (p/\alpha)^{\ell}}, \quad \text{for } \lambda \in \mathbb{Y}_{\ell}^{(p)}. \quad (3.1.4)$$

Probability $P_{u,p}(\lambda)$ is ≥ 0 , because, from (2.3.4), $J_{\lambda}^{(\alpha)}(1^p) > 0$, for $p \geq \lambda_1^{\top}$ and $= 0$ otherwise.

In particular, setting $\alpha = 1$ and using $J_{\lambda}^{(1)} = h^{\lambda} s_{\lambda}$ and $j_{\lambda}^{(1)} = (h^{\lambda})^2$, (3.1.3) leads to

$$P_{u,p}(\lambda) \Big|_{\alpha=1} = \frac{u^{|\lambda|} J_{\lambda}^{(1)}(1^p)}{e^{pu} j_{\lambda}^{(1)}} = \frac{u^{|\lambda|} s_{\lambda}(1^p)}{e^{pu} h^{\lambda}} = e^{-pu} \frac{(up)^{|\lambda|}}{|\lambda|!} P^{\ell,p}(\lambda) \Big|_{\alpha=1}, \quad \lambda \in \mathbb{Y}^{(p)}, \quad (3.1.5)$$

where by (2.1.2)

$$P^{\ell,p}(\lambda) \Big|_{\alpha=1} = \frac{f^{\lambda} s_{\lambda}(1^p)}{p^{|\lambda|}}, \quad \lambda \in \mathbb{Y}_{\ell}^{(p)}. \quad (3.1.6)$$

Probability (3.1.6) will be considered next.

$${}^7 \prod_{i \geq 1} (1 - x_i z)^{-1} = \exp \left(\sum_{\ell=1}^{\infty} \frac{z^{\ell}}{\ell} \sum_{i \geq 1} x_i^{\ell} \right).$$

3.2 Probability measures on the set $\mathbb{Y}_\ell$ of partitions of ℓ and random words

From (3.1.6), setting in this section $P^{\ell,p} := P^{\ell,p} \Big|_{\alpha=1}$,

$$\begin{aligned}
P^{\ell,p}(\lambda) &= \frac{f^\lambda s_\lambda(1^p)}{p^{|\lambda|}}, \quad \text{for } \lambda \in \mathbb{Y}_\ell^{(p)} \\
&= \frac{\ell!}{p^\ell} \Delta_p(h)^2 \frac{1}{\prod_1^{p-1} i! \prod_1^p h_i!}, \quad \text{by (2.1.2) and (2.1.3),} \\
&= \text{probability on Young diagrams } \lambda \in \mathbb{Y}_\ell^{(p)} \text{ coming from the} \\
&\quad \text{uniform distribution on } S_\ell^p, \text{ via the RSK correspondence,}
\end{aligned} \tag{3.2.1}$$

where $h := (h_1, \dots, h_p)$ with $h_i := p + \lambda_i - i$, and where

$$S_\ell^p := \{\text{words of length } \ell, \text{ built from an alphabet } \{1, \dots, p\}\},$$

with $|S_\ell^p| = p^\ell$. As already pointed out, this is a probability, firstly because of the connection with the word problem, secondly because of (3.1.5). As already pointed out, this probability was considered in [10, 11, 12, 50, 31] and also in the context of random words, by Tracy and Widom ([48]).

Proposition 3.1 *Given rectangular Young diagrams $\mu = (n-p)^p \supseteq \mu' = (n-q)^p$, with $p \leq q < n$, and $\ell \geq p(n-p)$, the expectation equals*

$$\begin{aligned}
&E^{\ell,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}} \right) \\
&= \frac{\ell!}{p^\ell} \prod_1^p \frac{(q-i)!}{(n-i)!} \sum_{\substack{\kappa \in \mathbb{Y}_{\ell-p(n-p)} \\ \kappa_1^\top \leq p}} \frac{1}{(h^\kappa)^2} \prod_{i=1}^p \frac{(p-i+1)_{\kappa_i} (q-i+1)_{\kappa_i}}{(n-i+1)_{\kappa_i}}. \tag{3.2.2}
\end{aligned}$$

Proof: From (2.2.1) with $\lambda \mapsto \lambda \setminus \mu'$, (so $n \mapsto q$, $p \mapsto p$) we have, upon setting $\kappa = \lambda \setminus \mu$,

$$\prod_{(i,j) \in \mu \setminus \mu'} h_{(i,j)}^\lambda = \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}} = h^{\mu \setminus \mu'} \prod_{i=1}^p \frac{(q-i+1)_{\kappa_i}}{(p-i+1)_{\kappa_i}}. \tag{3.2.3}$$

Then combining (3.2.3) and (2.2.3),

$$\frac{s_\lambda(1^p)}{h^\lambda} \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}} = \frac{h^{\mu \setminus \mu'}}{h^\mu} \frac{1}{(h^{\lambda \setminus \mu})^2} \prod_{i=1}^p \frac{(q-i+1)_{\kappa_i} (p-i+1)_{\kappa_i}}{(n-i+1)_{\kappa_i}}.$$

In particular, setting $\lambda = \mu$, and using (2.2.2),

$$\left. \frac{s_\lambda(1^p)}{h^\lambda} \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}} \right|_{\lambda=\mu} = \frac{h^{\mu \setminus \mu'}}{h^\mu} = \prod_1^p \frac{(q-i)!}{(n-i)!},$$

and therefore

$$\begin{aligned} \frac{s_\lambda(1^p)}{h^\lambda} \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}} &= \left(\frac{s_\lambda(1^p)}{h^\lambda} \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}} \right) \bigg|_{\lambda=\mu} \frac{1}{(h^{\lambda \setminus \mu})^2} \prod_{i=1}^p \frac{(p-i+1)_{\kappa_i} (q-i+1)_{\kappa_i}}{(n-i+1)_{\kappa_i}} \\ &= \prod_{i=1}^p \frac{(q-i)!}{(n-i)!} \frac{1}{(h^{\lambda \setminus \mu})^2} \prod_{i=1}^p \frac{(p-i+1)_{\kappa_i} (q-i+1)_{\kappa_i}}{(n-i+1)_{\kappa_i}}. \end{aligned} \quad (3.2.4)$$

Then, taking the expectation $E^{\ell,p}$ with regard to the probability measure $P^{\ell,p}$, defined in (3.2.1),

$$\begin{aligned} &E^{\ell,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{n-q < j \leq n-p} h_{(i,j)}^\lambda \right) \\ &= \sum_{\substack{\lambda \in \mathbb{Y}_\ell \\ \lambda \supseteq \mu}} \frac{f^\lambda s_\lambda(1^p)}{p^\ell} \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}} \\ &= \frac{\ell!}{p^\ell} \sum_{\substack{\lambda \in \mathbb{Y}_\ell \\ \lambda \supseteq \mu}} \frac{s_\lambda(1^p)}{h^\lambda} \frac{h^{\lambda \setminus \mu'}}{h^{\lambda \setminus \mu}}, \quad \text{using } f^\lambda = \frac{|\lambda|!}{h^\lambda} \text{ as in (2.1.2),} \\ &= \frac{\ell!}{p^\ell} \prod_1^p \frac{(q-i)!}{(n-i)!} \sum_{\substack{\lambda \in \mathbb{Y}_\ell \\ \lambda \supseteq \mu}} \frac{1}{(h^{\lambda \setminus \mu})^2} \prod_{i=1}^p \frac{(p-i+1)_{(\lambda \setminus \mu)_i} (q-i+1)_{(\lambda \setminus \mu)_i}}{(n-i+1)_{(\lambda \setminus \mu)_i}}, \\ &\hspace{25em} \text{using (3.2.4)} \end{aligned}$$

$$= \frac{\ell!}{p^\ell} \prod_1^p \frac{(q-i)!}{(n-i)!} \sum_{\substack{\kappa \in \mathbb{Y}_{\ell-p(n-p)} \\ \kappa_1^\top \leq p}} \frac{1}{(h^\kappa)^2} \prod_{i=1}^p \frac{(p-i+1)_{\kappa_i} (q-i+1)_{\kappa_i}}{(n-i+1)_{\kappa_i}}.$$

4 Expressing integrals as mathematical expectation on partitions and on random words

4.1 Expressing an integral on $Gr(p, \mathbb{F}^n)$ as a mathematical expectation on partitions

Remembering the probability (3.1.3)

$$P_{x,p}(\lambda) = e^{-\beta p x} \frac{x^{|\lambda|} J_{\lambda}^{(1/\beta)}(1^p)}{j_{\lambda}^{(1/\beta)}}, \quad \lambda \in \mathbb{Y}, \quad (4.1.1)$$

on partitions $\mathbb{Y}$, with support on $\lambda_1^{\top} \leq p$, the following statement holds:

Theorem 4.1 *For fixed $p \leq q \leq n/2$, the following holds ($\beta = 1/2, 1, 2$)*

$$c^{-1} \int_{Gr(p, \mathbb{F}^n)} e^{x \operatorname{Tr}(I+Z^{\dagger}Z)^{-1}} \det(Z^{\dagger}Z)^{-\beta(q-p)} d\mu(Z) \quad (4.1.2)$$

$$= {}_2F_1^{(1/\beta)}(\beta p, \beta q; \beta n; y) \Big|_{\sum_{\ell} y_{\ell}^i = \frac{x}{\beta} \delta_{1i}} \quad (4.1.3)$$

$$= e^{\beta p x} E_{x,p} \left(\frac{1}{\beta^{|\lambda|}} \frac{J_{\lambda}^{(1/\beta)}(1^q)}{J_{\lambda}^{(1/\beta)}(1^n)} \right) \quad (4.1.4)$$

where $c := c_{n,q,p}^{(\beta)}$ is as in (0.0.20).

Proof: For a symmetric function $f(z_1, \dots, z_p)$, define the integral, depending on β ,

$$\langle f \rangle_{\beta} := \int_{[0,1]^p} f(z_1, \dots, z_p) |\Delta(z)|^{2\beta} \prod_{i=1}^p z_i^{\beta(q-p+1)-1} (1-z_i)^{\beta(n-p-q+1)-1} dz_i. \quad (4.1.5)$$

Kaneko's formula (2.3.13) will be used for $a = \beta(n-q)$ and $b = \beta q$ in the sequence of identities below; the inequalities $p \leq q \leq n/2$ imply $q-p+1, n-p-q+1 \geq 1$, and so the integral (4.1.5) above makes sense, and we first apply (1.0.3) in the following sequence of identities:

$$\begin{aligned}
& \int_{Gr(p, \mathbb{F}^n)} e^{x \operatorname{Tr}(I+Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-\beta(q-p)} d\mu(Z) \\
&= \left\langle \prod_{i=1}^p e^{xz_i} \right\rangle_\beta \\
&= \sum_0^\infty x^\ell \left\langle \frac{\left(\sum_1^p z_i\right)^\ell}{\ell!} \right\rangle_\beta \\
&= \sum_0^\infty x^\ell \left\langle \frac{1}{\beta^\ell} \sum_{|\lambda|=\ell} \frac{J_\lambda^{(1/\beta)}(z)}{j_\lambda^{(1/\beta)}} \right\rangle_\beta, \text{ using (2.3.6)} \\
&\stackrel{*}{=} \prod_{i=1}^p \frac{\Gamma(i\beta+1)\Gamma(\beta(n-q-i+1))}{\Gamma(\beta+1)} \\
&\quad \sum_{\ell=0}^\infty \left(\frac{x}{\beta}\right)^\ell \sum_{|\lambda|=\ell} \frac{J_\lambda^{(1/\beta)}(1^p)}{j_\lambda^{(1/\beta)}} \prod_{i=1}^p \frac{\Gamma(\lambda_i + \beta(q-i+1))}{\Gamma(\lambda_i + \beta(n-i+1))}, \\
&\quad \text{using Kaneko's formula (2.3.13),} \\
&= \prod_{i=1}^p \frac{\Gamma(i\beta+1)\Gamma(\beta(n-q-i+1))}{\Gamma(\beta+1)} e^{\beta p x} \\
&\quad \sum_{\lambda \in \mathbb{Y}} \frac{1}{\beta^{|\lambda|}} P_{x,p}(\lambda) \prod_{i=1}^p \frac{\Gamma(\lambda_i + \beta(q-i+1))}{\Gamma(\lambda_i + \beta(n-i+1))}, \quad \begin{array}{l} \text{using the definition} \\ \text{(4.1.1) of probability,} \\ \text{with support on } \mathbb{Y}^{(p)} \end{array} \\
&= c_{n,q,p}^{(\beta)} e^{\beta p x} \sum_{\lambda \in \mathbb{Y}} P_{x,p}(\lambda) \frac{J_\lambda^{(1/\beta)}(1^q)}{\beta^{|\lambda|} J_\lambda^{(1/\beta)}(1^n)}, \quad \begin{array}{l} \text{using formula (2.3.4) and} \\ \text{the value (0.0.20) of } c_{n,q,p}^{(\beta)}, \end{array} \\
&= c_{n,q,p}^{(\beta)} e^{\beta p x} E_{x,p} \left(\frac{1}{\beta^{|\lambda|}} \frac{J_\lambda^{(1/\beta)}(1^q)}{J_\lambda^{(1/\beta)}(1^n)} \right), \tag{4.1.6}
\end{aligned}$$

yielding (4.1.4).

Finally, looking at the expression to the right of $\stackrel{*}{=}$, we find, using (3.1.2) and (2.3.4), to be precise,

$$J_\lambda^{(1/\beta)}(y) \Big|_{\sum_\ell y_\ell^i = \frac{x}{\beta} \delta_{1i}} = \left(\frac{x}{\beta}\right)^{|\lambda|} \quad \text{and} \quad J_\lambda^{(1/\beta)}(1^p) = \left(\frac{1}{\beta}\right)^{|\lambda|} (p\beta)_\lambda^{(1/\beta)},$$

the following

$$\begin{aligned}
\left\langle \prod_{i=1}^p e^{xz_i} \right\rangle_{\beta} &= c_{n,q,p}^{(\beta)} \sum_{\lambda \in \mathbb{Y}} \frac{(\beta p)_{\lambda} (\beta q)_{\lambda}}{(\beta n)_{\lambda}} \frac{1}{\beta^{|\lambda|}} \frac{J_{\lambda}^{(1/\beta)}(y)}{j_{\lambda}^{(1/\beta)}} \Big|_{\sum_{\ell} y_{\ell}^i = \frac{x}{\beta} \delta_{1i}}, \quad \text{by (2.3.4),} \\
&= c_{n,q,p}^{(\beta)} {}_2F_1^{(1/\beta)}(\beta p, \beta q; \beta n; y) \Big|_{\sum_{\ell} y_{\ell}^i = \frac{x}{\beta} \delta_{1i}}, \quad \text{by (2.3.10),}
\end{aligned}$$

thus ending the proof of Theorem 4.1. ■

Remark: Identity (4.1.3) is also an immediate consequence of Kaneko's formula.

4.2 Expressing an integral over $Gr(p, \mathbb{C}^n)$ as a mathematical expectation on partitions and random words

We now specialize the previous section to $\mathbb{F} = \mathbb{C}$. For fixed integer $p \geq 1$ and $\beta = 1$, recall the probability (3.1.5), with support in $\mathbb{Y}^{(p)}$,

$$P_{x,p}(\lambda) = \frac{x^{|\lambda|} s_{\lambda}(1^p)}{e^{px} h^{\lambda}} = e^{-px} \frac{(xp)^{|\lambda|}}{|\lambda|!} P^{\ell,p}(\lambda), \quad \lambda \in \mathbb{Y}^{(p)}, \quad (4.2.1)$$

and the probability (3.1.6) on $\mathbb{Y}_{\ell}^{(p)}$ coming from the uniform distribution on S_{ℓ}^p , via the RSK correspondence,

$$P^{\ell,p}(\lambda) = \frac{f^{\lambda} s_{\lambda}(1^p)}{p^{|\lambda|}}, \quad \lambda \in \mathbb{Y}_{\ell}^{(p)}. \quad (4.2.2)$$

For integer $0 \leq p \leq n/2$, consider the fixed rectangular Young diagram $\mu = (n-p)^p$.

Theorem 4.2 *For fixed $p \leq q \leq n/2$,*

$$\begin{aligned}
& \int_{Gr(p, \mathbb{C}^n)} e^{x \operatorname{Tr}(I+Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-(q-p)} d\mu(Z) \\
&= \left\{ \begin{aligned} & \frac{\tilde{c}_{n,q,p}}{x^{(n-p)p}} e^{px} E_{x,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^\lambda \right) \\ & \frac{\tilde{c}_{n,q,p}}{x^{(n-p)p}} \sum_{\ell \geq p(n-p)} \frac{(px)^\ell}{\ell!} E^{\ell,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^\lambda \right), \\ & c_{n,q,p}^{(1)} \sum_{r \geq 0} x^r \sum_{\substack{\kappa \in \mathbb{Y}_r \\ \kappa_1^\top \leq p}} \frac{1}{(h^\kappa)^2} \frac{(p)_\kappa (q)_\kappa}{(n)_\kappa} \\ & c_{n,q,p}^{(1)} {}_2F_1^{(1)}(p, q; n; y) \Big|_{\sum_\ell y_\ell^i = \delta_{1i} x} \end{aligned} \right. \quad (4.2.3)
\end{aligned}$$

In particular, for $p = q$, the integral above has two different formulations, as a probability or as a generating function of probabilities,

$$\begin{aligned}
& \int_{Gr(p, \mathbb{C}^n)} e^{x \operatorname{Tr}(I+Z^\dagger Z)^{-1}} d\mu(Z) \\
&= \left\{ \begin{aligned} & \frac{\tilde{c}_{n,p,p}}{x^{(n-p)p}} e^{px} P_{x,p}(\lambda \supseteq \mu) \\ & \frac{\tilde{c}_{n,p,p}}{x^{(n-p)p}} \sum_{\ell \geq p(n-p)} \frac{(px)^\ell}{\ell!} P^{\ell,p}(\lambda \supseteq \mu), \\ & \frac{\tilde{c}_{n,p,p}}{x^{(n-p)p}} \sum_{\ell \geq p(n-p)} \frac{(px)^\ell}{\ell!} P^{\ell,p} \left(\pi \in S_\ell^p \mid \begin{array}{l} d_1(\pi) = p \text{ and} \\ i_{p-1}(\pi) \leq \ell - n + p \end{array} \right), \\ & c_{n,p,p}^{(1)} \sum_{r \geq 0} x^r \sum_{\substack{\kappa \in \mathbb{Y}_r \\ \kappa_1^\top \leq p}} \frac{1}{(h^\kappa)^2} \frac{((p)_\kappa)^2}{(n)_\kappa} \\ & c_{n,p,p}^{(1)} {}_2F_1^{(1)}(p, p; n; y) \Big|_{\sum_\ell y_\ell^i = \delta_{1i} x} \end{aligned} \right.
\end{aligned}$$

$$(4.2.4)$$

where $P^{\ell,p}$ denotes the uniform distribution on the set S_ℓ^p of words of length ℓ from an alphabet $1, \dots, p$, and

$d_1(\pi)$ = length of longest strictly decreasing subsequence

$i_k(\pi)$ = length of the longest union of k disjoint weakly increasing subsequences.

Remark: The constant $c_{n,q,p}^{(1)}$ in (4.2.2) and (4.2.3) is the same as $c_{n,q,p}^{(\beta)}$ for $\beta = 1$ (see (0.0.20)) and $\tilde{c}_{n,q,p}$ is a new constant:

$$c_{n,q,p}^{(1)} = \prod_{j=1}^p \binom{n-j}{n-q-j, q-j, j}^{-1} \quad \tilde{c}_{n,q,p} = \prod_{i=1}^p i! (n-q-i)! \quad (4.2.5)$$

From the fourth expression of (4.2.2), it follows readily that, near $x = 0$,

$$\begin{aligned} & (c_{n,q,p}^{(1)})^{-1} \int_{Gr(p, \mathbb{C}^n)} e^{x \operatorname{Tr}(I+Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-(q-p)} d\mu(Z) \\ &= 1 + \frac{p}{n} x + \frac{pq}{4n} \left(\frac{(p+1)(q+1)}{n+1} + \frac{(p-1)(q-1)}{n-1} \right) x^2 + \dots \end{aligned} \quad (4.2.6)$$

Proof: From the last identity in (2.1.2) with $m = p$, it follows that, since $\lambda_1^\top \leq p$,

$$\frac{h^{\lambda+(q-p)p}}{h^{\lambda+(n-p)p}} = \prod_{i=1}^p \frac{(q+\lambda_i-i)!}{(n+\lambda_i-i)!} \quad (4.2.7)$$

For a partition λ such that $\lambda_1^\top \leq p$ and for $\lambda' = \lambda + k^p$, with arbitrary integer $k \geq 0$, we have $s_\lambda(1^p) = s_{\lambda'}(1^p)$, using the last identity (2.1.3). Using these facts, we have, continuing from (4.1.6),

$$\begin{aligned} & \int_{Gr(p, \mathbb{C}^n)} e^{x \operatorname{Tr}(I+Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{p-q} d\mu(Z) \\ &= \int_{[0,1]^p} \Delta(z)^2 \prod_{i=1}^p e^{xz_i} z_i^{q-p} (1-z_i)^{n-p-q} dz_i \\ &= \tilde{c}_{n,q,p} \sum_{\substack{\lambda \in \mathbb{Y} \\ \lambda_1^\top \leq p}} x^{|\lambda|} \frac{s_\lambda(1^p)}{h^\lambda} \prod_{i=1}^p \frac{(q+\lambda_i-i)!}{(n+\lambda_i-i)!}, \quad \text{by the 4th identity of (4.1.6) and } J_\lambda^{(1)} = h^\lambda s_\lambda \text{ and } j_\lambda^{(1)} = (h^\lambda)^2 \\ &= \tilde{c}_{n,q,p} \sum_{\lambda \in \mathbb{Y}} x^{|\lambda|} \frac{s_\lambda^p}{h^\lambda} \frac{h^{\lambda+(q-p)p}}{h^{\lambda+(n-p)p}}, \quad \text{using (4.2.7)} \end{aligned}$$

$$\begin{aligned}
& \stackrel{*}{=} \tilde{c}_{n,q,p} \sum_{\substack{\lambda' \in \mathbb{Y} \\ \lambda' \supseteq \mu}} x^{|\lambda'| - (n-p)p} \frac{s_{\lambda'}(1^p)}{h^{\lambda'}} \frac{h^{\lambda' - (n-q)p}}{h^{\lambda' - (n-p)p}}, \quad \begin{array}{l} \text{setting } \lambda' = \lambda + (n-p)p \\ \text{and using } s_{\lambda}(1^p) = s_{\lambda'}(1^p) \end{array} \\
& = \frac{\tilde{c}_{n,q,p} e^{px}}{x^{(n-p)p}} \sum_{\substack{\lambda' \in \mathbb{Y} \\ \lambda' \supseteq \mu}} P_{x,p}(\lambda') \frac{h^{\lambda' - (n-q)p}}{h^{\lambda' - (n-p)p}}, \quad \begin{array}{l} \text{using } P_{x,p} \text{ defined in (4.2.1),} \\ \text{and } |\lambda'| = |\lambda| + (n-p)p, \end{array} \\
& = \frac{\tilde{c}_{n,q,p} e^{px}}{x^{(n-p)p}} E_{x,p} \left(I_{\lambda \supseteq \mu}(\lambda) \prod_{n-q < j \leq n-p} h_{(i,j)}^{\lambda} \right). \tag{4.2.8}
\end{aligned}$$

The last equality in (4.2.3) follows from (4.1.3), while the second to the last one follows from (2.3.12).

Finally, to prove the second formula on the right hand side of (4.2.3), start with equality $\stackrel{*}{=}$ in (4.2.8), omitting $\tilde{c}_{n,q,p}$, and replacing λ' by λ ,

$$\begin{aligned}
& \frac{1}{x^{p(n-p)}} \sum_{\substack{\lambda \in \mathbb{Y} \\ \lambda \supseteq \mu}} x^{|\lambda|} \frac{s_{\lambda}(1^p)}{h^{\lambda}} \frac{h^{\lambda - (n-q)p}}{h^{\lambda - (n-p)p}}, \quad \text{for fixed } \mu = (n-p)^p \\
& = \frac{1}{x^{p(n-p)}} \sum_{\substack{\lambda \in \mathbb{Y} \\ \lambda \supseteq \mu}} \frac{(px)^{|\lambda|}}{|\lambda|!} \frac{|\lambda|!}{h^{\lambda}} \frac{s_{\lambda}(1^p)}{p^{|\lambda|}} \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^{\lambda} \\
& = \frac{1}{x^{p(n-p)}} \sum_{\ell \geq p(n-p)} \frac{(px)^{\ell}}{\ell!} \sum_{\substack{\lambda \in \mathbb{Y}_{\ell} \\ \lambda \supseteq \mu}} \frac{f^{\lambda} s_{\lambda}(1^p)}{p^{\ell}} \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^{\lambda}, \quad \text{using (2.1.2),} \\
& = \frac{1}{x^{p(n-p)}} \sum_{\ell \geq p(n-p)} \frac{(px)^{\ell}}{\ell!} E^{\ell,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^{\lambda} \right).
\end{aligned}$$

In particular, setting $q = p$, the latter equals

$$\begin{aligned}
& \frac{1}{x^{p(n-p)}} \sum_{\ell \geq p(n-p)} \frac{(px)^{\ell}}{\ell!} P^{\ell,p}(\lambda \supseteq \mu) \\
& = \frac{1}{x^{p(n-p)}} \sum_{\ell \geq p(n-p)} \frac{(px)^{\ell}}{\ell!} P^{\ell,p}(\lambda \supseteq \mu, \lambda_1^{\top} = p) \\
& = \frac{1}{x^{p(n-p)}} \sum_{\ell \geq p(n-p)} \frac{(px)^{\ell}}{\ell!} P^{\ell,p}(\pi \in S_{\ell}^p \mid d_1(\pi) = p, \ i_{p-1}(\pi) \leq \ell - n + p).
\end{aligned}$$

To see the last two equalities, one proceeds as follows. From $\lambda \supseteq \mu = (n-p)^p$ and $P^{\ell,p}(\lambda) = 0$ for $\lambda_1^\top > p$, it follows that $\lambda_1^\top = p$ and by Greene's theorem (see Sagan [41], p. 110), $d_1(\pi) = \lambda_1^\top = p$. Since also by Greene, $i_k(\pi) = \sum_1^k \lambda_i$, and, in particular $i_p(\pi) = \sum_1^p \lambda_i = \ell$, we have

$$\lambda_p = i_p(\pi) - i_{p-1}(\pi) = \ell - i_{p-1}(\pi).$$

From $\lambda \supseteq \mu = (n-p)^p$, it also follows that $\lambda_p \geq n-p$, and thus $i_{p-1}(\pi) \leq \ell - n + p$. Conversely, if $d_1(\pi) = \lambda_1^\top = p$, and $i_{p-1}(\pi) \leq \ell - n + p$, then $\sum_1^{p-1} \lambda_i = i_{p-1}(\pi) \leq \sum_1^p \lambda_i - n + p$; hence $\lambda_p \geq n-p$, and so $\lambda \supseteq \mu$. ■

5 Testing Statistical Independence of Gaussian Populations

The statistical facts, used in this paper and summarized in this section, are due to James [25] and Constantine [13]; see also Muirhead [35].

5.1 The Wishart distribution

Let the $p \times n$ matrix X , with $n \geq p$ and n identically distributed independent columns, have the normal distribution

$$(\det 2\pi\Sigma)^{-n/2} e^{-\frac{1}{2} \text{Tr} \Sigma^{-1}(X-M)(X-M)^\top}. \quad (5.1.1)$$

Then the $p \times p$ matrix $S = X X^\top$ has the *non-central Wishart distribution with n degrees of freedom*, with $p \times p$ covariance matrix Σ and non-centrality matrix $\Omega = \frac{1}{2} M M^\top \Sigma^{-1}$, namely

$$\Gamma_p(n/2)^{-1} (\det 2\Sigma)^{-n/2} e^{-\text{Tr}(\Omega + \frac{1}{2}\Sigma^{-1}S)} (\det S)^{\frac{1}{2}(n-p-1)} {}_0F_1\left(\frac{n}{2}; \frac{1}{2}\Sigma^{-1}\Omega S\right),$$

where $S > 0$, where Γ_m is the multivariate Gamma function and where⁸(for the definition of C_λ , see (2.3.7))

$${}_0F_1\left(\frac{n}{2}; \frac{1}{2}\Sigma^{-1}\Omega S\right) = \sum_{\lambda \in \mathbb{Y}} \frac{C_\lambda \left(\frac{1}{2}\Sigma^{-1}\Omega S\right)}{\left(\frac{n}{2}\right)_\lambda |\lambda|!} = \sum_{\lambda \in \mathbb{Y}} \frac{2^{|\lambda|} J_\lambda^{(2)} \left(\frac{1}{2}\Sigma^{-1}\Omega S\right)}{\left(\frac{n}{2}\right)_\lambda j_k^{(2)}}.$$

⁸Here $(a)_\lambda := \prod_i (a + \beta(1-i))_{\lambda_i}$ for $\beta = 1/2$, with $(x)_n := x(x+1) \dots (x+n-1)$, $x_0 = 1$.

When $M = 0$, we find the (*central*) *Wishart distribution* $W_p(n, \Sigma)$, with $p \leq n$, for the $p \times p$ matrix $S = X X^\top$:

$$\Gamma_p(n/2)^{-1} (\det 2\Sigma)^{-n/2} e^{-\frac{1}{2} \text{Tr} \Sigma^{-1} S} (\det S)^{\frac{1}{2}(n-p-1)} \prod_{1 \leq i \leq j \leq p} dS_{ij}. \quad (5.1.2)$$

5.2 The canonical correlation coefficients

In testing the statistical independence of two Gaussian populations, one needs to know the distribution of *canonical correlation coefficients*. To set up the problem, consider $p + q$ normally distributed random variables $(X_1, \dots, X_p)^\top$ and $(Y_1, \dots, Y_q)^\top$ ($p \leq q$) with mean zero and covariance matrix

$$\begin{array}{c} \xleftrightarrow{p} \quad \xleftrightarrow{q} \\ \text{cov} \left(\begin{array}{c} X \\ Y \end{array} \right) := \Sigma = \left(\begin{array}{cc} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{12}^\top & \Sigma_{22} \end{array} \right) \begin{array}{c} \updownarrow p \\ \updownarrow q \end{array} \end{array}$$

The method proposed by Hotelling [24] is to find linear transformations $U = L_1 X$ and $V = L_2 Y$ of X and Y having the property that the correlation between the first components U_1 and V_1 of the vectors U and V is maximal subject to the condition that $\text{Var } U_1 = \text{Var } V_1 = 1$; moreover, one requires the second components U_2 and V_2 to have maximal correlation subjected to

$$\begin{cases} \text{(i)} & \text{Var } U_2 = \text{Var } V_2 = 1 \\ \text{(ii)} & U_2 \text{ and } V_2 \text{ are uncorrelated with both } U_1 \text{ and } V_1, \end{cases}$$

etc ...

Then there exist $O_p \in O(p)$, $O_q \in O(q)$ such that

$$\Sigma_{11}^{-1/2} \Sigma_{12} \Sigma_{22}^{-1/2} = O_p^\top P O_q$$

where P has the following form:

$$\left\{ \begin{array}{l}
P = \left(\begin{array}{cccc|c}
\rho_1 & & & & \\
& \ddots & & & \\
& & O & & \\
& & \rho_k & & \\
& & \rho_{k+1} & & \\
O & & & \ddots & \\
& & & & \rho_p
\end{array} \right) \begin{array}{l} \xleftrightarrow{q} \\ \updownarrow p, \quad k = \text{rank } \Sigma_{12}, \\ \xleftrightarrow{p} \end{array} \\
1 \geq \rho_1 \geq \rho_2 \geq \dots \geq \rho_k > 0, \quad \rho_{k+1} = \dots = \rho_p = 0 \quad (\text{canonical correlation coefficients}), \\
\rho_i \text{ are solutions } (\geq 0) \text{ of } \det(\Sigma_{11}^{-1} \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{12}^\top - \rho^2 I) = 0.
\end{array} \right.$$

Then the covariance matrix of the vectors

$$U = L_1 X := O_p \Sigma_{11}^{-1/2} X \quad \text{and} \quad V = L_2 Y := O_q \Sigma_{22}^{-1/2} Y$$

has the canonical form ($\det \Sigma_{can} = \prod_1^p (1 - \rho_i^2)$)

$$\text{cov} \begin{pmatrix} U \\ V \end{pmatrix} = \Sigma_{can} = \begin{pmatrix} I_p & P \\ P^\top & I_q \end{pmatrix},$$

with

$$\text{spectrum } \Sigma_{can} = \underbrace{1, \dots, 1}_{q-p}, 1 - \rho_1, 1 + \rho_1, \dots, 1 - \rho_p, 1 + \rho_p$$

and inverse

$$\Sigma_{can}^{-1} = \frac{1}{\prod_1^p (1 - \rho_i^2)^2} \begin{pmatrix} I_p & -P \\ -P^\top & I_q \end{pmatrix}.$$

5.3 Distribution of the sample canonical correlations

From here on, we may take $\Sigma = \Sigma_{can}$. The n ($n \geq p + q$) independent samples $(x_{11}, \dots, x_{1p}, y_{11}, \dots, y_{1q})^\top, \dots, (x_{n1}, \dots, x_{np}, y_{n1}, \dots, y_{nq})^\top$, arising from observing $\begin{pmatrix} X \\ Y \end{pmatrix}$ lead to a matrix $\begin{pmatrix} x \\ y \end{pmatrix}$ of size $(p + q, n)$, having the normal distribution [35] (p. 79 and p. 539)

$$\begin{aligned}
& (2\pi)^{-n(p+q)/2} (\det \Sigma)^{-n/2} \exp -\frac{1}{2} \text{Tr} \begin{pmatrix} x^\top & y^\top \end{pmatrix} \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{12}^\top & \Sigma_{22} \end{pmatrix}^{-1} \begin{pmatrix} x \\ y \end{pmatrix} \\
& = (2\pi)^{-n(p+q)/2} (\det \Sigma)^{-n/2} e^{-\frac{1}{2} \text{Tr} \begin{pmatrix} x^\top (\Sigma^{-1})_{11} x + y^\top (\Sigma^{-1})_{22} y + 2y^\top (\Sigma^{-1})_{12}^\top x \end{pmatrix}}
\end{aligned}$$

The conditional distribution of $p \times n$ matrix x given the $q \times n$ matrix y is also normal:

$$(\det 2\pi\Omega)^{-n/2} e^{-\frac{1}{2} \text{Tr} \Omega^{-1}(x-Py)(x-Py)^\top} \quad (5.3.1)$$

with

$$\begin{aligned}
\Omega &= \Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21} = \text{diag}(1 - \rho_1^2, \dots, 1 - \rho_p^2) \\
P &= \Sigma_{12} \Sigma_{22}^{-1}.
\end{aligned}$$

Then the maximum likelihood estimates r_i of the ρ_i satisfy the determinantal equation

$$\det(S_{11}^{-1} S_{12} S_{22}^{-1} S_{12}^\top - r^2 I) = 0, \quad (5.3.2)$$

corresponding to

$$S = \begin{pmatrix} S_{11} & S_{12} \\ S_{12}^\top & S_{22} \end{pmatrix} := \begin{pmatrix} xx^\top & xy^\top \\ yx^\top & yy^\top \end{pmatrix},$$

where S_{ij} are the associated submatrices of the *sample* covariance matrix S .

Remark: The r_i can also be viewed as $r_i = \cos \theta_i$, where the $\theta_1, \dots, \theta_p$ are the *critical* angles between two planes in $\mathbb{R}^n$:

- (i) a p -dimensional plane = $\text{span} \{(x_{11}, \dots, x_{n1}), \dots, (x_{1p}, \dots, x_{np})\}$
- (ii) a q -dimensional plane = $\text{span} \{(y_{11}, \dots, y_{n1})^\top, \dots, (y_{1q}, \dots, y_{nq})^\top\}$.

As we shall see, $z_i = r_i^2 = \cos^2 \theta_i$ are the precise variables z_i appearing in section 1.

Since the (q, n) -matrix y has $\text{rank}(y) = q$, there exists a matrix $H_n \in O(n)$ such that $yH_n = (y_1 \mid O)$; therefore acting on x with H_n leads to

$$\begin{aligned}
& \begin{matrix} \xleftrightarrow{q} & \xleftrightarrow{n-q} \end{matrix} & \begin{matrix} \xleftrightarrow{q} & \xleftrightarrow{n-q} \end{matrix} \\
yH = (y_1 \mid O) \uparrow q, & \quad xH_n = (u \mid v) \uparrow p. \quad (5.3.3)
\end{aligned}$$

With this in mind,

$$\begin{aligned}
& S_{12}S_{22}^{-1}S_{12}^\top - r^2S_{11} \\
&= xy^\top(yy^\top)^{-1}yx^\top - r^2xx^\top \\
&= xH(yH)^\top(yH(yH)^\top)^{-1}yH(xH)^\top - r^2(xH)(xH)^\top \\
&= (u \mid v) \begin{pmatrix} y_1^\top \\ O \end{pmatrix} \left((y_1 \mid O) \begin{pmatrix} y_1^\top \\ O \end{pmatrix} \right)^{-1} (y_1 \mid O) \begin{pmatrix} u^\top \\ v^\top \end{pmatrix} - r^2(u \mid v) \begin{pmatrix} u^\top \\ v^\top \end{pmatrix} \\
&= (u \mid v) \begin{pmatrix} I_q & O \\ O & 0_{n-q} \end{pmatrix} \begin{pmatrix} u^\top \\ v^\top \end{pmatrix} - r^2(u \mid v) \begin{pmatrix} u^\top \\ v^\top \end{pmatrix} \\
&= uu^\top - r^2(uu^\top + vv^\top),
\end{aligned}$$

and so the equation (5.3.2) for the r_i can be rewritten

$$\det(uu^\top - r^2(uu^\top + vv^\top)) = 0. \quad (5.3.4)$$

Then setting the forms (5.3.3) of x and y in the conditional distribution (5.3.1) of x given y , one computes the following, setting $H := H_n$,

$$\begin{aligned}
& \text{Tr } \Omega^{-1}(x - Py)(x - Py)^\top \\
&= \text{Tr } \Omega^{-1}(xH - PyH)(xH - PyH)^\top \\
&= \text{Tr } \Omega^{-1}((u \mid v) - P(y_1 \mid O))((u \mid v) - P(y_1 \mid O))^\top \\
&= \text{Tr } \Omega^{-1}(u - Py_1)(u - Py_1)^\top + \text{Tr } \Omega^{-1}vv^\top; \quad \Omega = \text{diag}(1 - \rho_1^2, \dots, 1 - \rho_p^2);
\end{aligned}$$

this establishes the independence of the normal distributions u and v , given the matrix y , with

$$u \equiv N(Py_1, \Omega), \quad v \equiv N(O, \Omega). \quad P = \text{diag}(\rho_1, \dots, \rho_p).$$

Hence $uu^\top$ and $vv^\top$ are conditionally independent and both Wishart distributed; to be precise:

- The $p \times p$ matrices $vv^\top$ are Wishart distributed, given y , with $n - q$ degrees of freedom and covariance Ω ;
- The $p \times p$ matrices $uu^\top$ are non-centrally Wishart distributed, given y , with q degrees of freedom, with covariance Ω and with non-centrality matrix

$$\frac{1}{2}Py_1y_1^\top P^\top \Omega^{-1}.$$

- The marginal distribution of the $q \times q$ matrices $yy^\top$ are Wishart distributed, with n degrees of freedom and covariance I_q , because the marginal distribution of y is normal with covariance I_q .

To summarize, given the matrix y , the sample canonical correlation coefficients $r_1^2 > \dots > r_p^2$ are the roots of

$$\begin{aligned} (r_1^2 > \dots > r_p^2) &= \text{roots of } \det(xy^\top(yy^\top)^{-1}yx^\top - r^2xx^\top) = 0 \\ &= \text{roots of } \det(uu^\top - r^2(uu^\top + vv^\top)) = 0 \\ &= \text{roots of } \det(uu^\top(uu^\top + vv^\top)^{-1} - r^2I) = 0. \end{aligned}$$

Then one shows that, knowing $uu^\top$ and $vv^\top$ are Wishart and conditionally independent, the conditional distribution of $r_1^2 > \dots > r_p^2$, given the matrix y is given by

$$\begin{aligned} \pi^{p^2/2} c_{n,p,q} e^{-\frac{1}{2} \text{Tr } Pyy^\top P^\top \Omega^{-1}} \Delta_p(r^2) \prod_{i=1}^p (r_i^2)^{\frac{1}{2}(q-p-1)} (1 - r_i^2)^{\frac{1}{2}(n-q-p-1)} \\ \sum_{\lambda \in \mathbb{Y}} \frac{(n/2)_\lambda C_\lambda(\frac{1}{2} Pyy^\top P^\top \Omega^{-1})}{(q/2)_\lambda C_\lambda(I_p) |\lambda|!} C_\lambda(R^2), \end{aligned}$$

where⁹

$$R^2 = \text{diag}(r_1^2, \dots, r_p^2), \quad c_{n,p,q} = \frac{\Gamma_p(n/2)}{\Gamma_p(q/2) \Gamma_p((n-q)/2) \Gamma_p(p/2)}.$$

By taking the expectation with regard to y or, what is the same, by integrating over the matrix $yy^\top$, which is Wishart distributed, we obtain:

Theorem 5.1 *Let $X_1, \dots, X_p, Y_1, \dots, Y_q$ ($p \leq q$) be normally distributed random variables with zero means and covariance matrix $\Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix}$. If $\rho_1^2, \dots, \rho_p^2$ are the roots of $\det(\Sigma_{11}^{-1} \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{12}^\top - \rho^2 I) = 0$, then the maximum likelihood estimates $r_1^2, \dots, r_p^2$ from a sample of size n ($n \geq p + q$) are given by the roots of*

$$\det(xy^\top(yy^\top)^{-1}yx^\top - r^2xx^\top) = 0.$$

Setting

$$Z := \text{diag}(z_1, \dots, z_p) = \text{diag}(r_1^2, \dots, r_p^2) \quad \text{and} \quad P^2 := \text{diag}(\rho_1^2, \dots, \rho_p^2),$$

⁹ $c_{n,p,q}$ is a different constant from (0.0.20).

the $z_i = r_i^2$ have the following density

$$\pi^{-p^2} c_{n,p,q} \Delta_p(z) \prod_{i=1}^p z_i^{(q-p-1)/2} (1-z_i)^{(n-q-p-1)/2} dz_i \cdot \left(\prod_{i=1}^p (1-\rho_i^2)^{n/2} \right) \sum_{\lambda \in \mathbb{Y}} \frac{\left(\frac{n}{2}\right)_\lambda \left(\frac{n}{2}\right)_\lambda}{\left(\frac{q}{2}\right)_\lambda} \frac{C_\lambda(Z) C_\lambda(P^2)}{C_\lambda(1^p) |\lambda|!}. \quad (5.3.5)$$

Corollary 5.2 *If $\rho_1^2 = \dots = \rho_p^2 = 0$, then the joint density of the $z_i = r_i^2$ is given by the density appearing in the integral of Theorem 1.1, namely*

$$\pi^{p^2/2} c_{n,p,q} \Delta_p(z) \prod_{i=1}^p z_i^{(q-p-1)/2} (1-z_i)^{(n-q-p-1)/2} dz_i. \quad (5.3.6)$$

Remark: As was shown here, the normal distribution over $\mathbb{R}$ leads to the density (5.3.6) for the z_i , which corresponds to the case $\beta = 1/2$ for (1.0.3). Starting with normal distributions over $\mathbb{C}$ and $\mathbb{H}$ leads, in a similar way, to integrals (1.0.3) for the cases $\beta = 1$ and 2.

6 Differential equations for the Grassmannian integrals and the hypergeometric functions

6.1 Differential equations for the Grassmannian integral

Theorem 6.1 shows that the integral over the Grassmannian $Gr(p, \mathbb{F}^n)$ satisfies Painlevé-like differential equations; for $\beta = 1$, this equation is the Painlevé V equation with a specific boundary condition (Theorem 6.2).

Theorem 6.1 *The following holds for the integral*

$$I_p(y) = \int_{Gr(p, \mathbb{F}^n)} e^{y \operatorname{Tr}(I+Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-\beta(q-p)} d\mu(Z) = c' \exp \int_0^x H(y) dy, \quad (6.1.1)$$

where $H(y) = \frac{d}{dy} \log I_p(y)$ satisfies the differential equation ($H' := \frac{dH}{dy}$ and remember $\delta_1^\beta = 1$ for $\beta = 1$ and $= 0$ otherwise):

$$4 \left(y^3 H''' + 6y^3 H'^2 + (1 + \delta_1^\beta)(2y^2 H'' + 4y^2 H H' + y H^2) \right) - y P_0 H' + P_1 H + P_2$$

$$= \begin{cases} 0, & \text{for } \beta = 1, \quad (\text{Painlevé V}) \\ \frac{3}{16} \frac{p(p-1)}{(p+1)(p+2)} y^3 \frac{I_{p-2} I_{p+2}}{I_p^2}, & \text{for } \beta = 1/2, \\ \frac{3}{16^2} \frac{p}{p+1} y^3 \frac{I_{p-1} I_{p+1}}{I_p^2}, & \text{for } \beta = 2, \end{cases} \quad (6.1.2)$$

where

	P_0	P_1	P_2
$\beta = 1$	$4y^2 - 8sy + 4n^2 - 8$	$4(sy - n^2)$	$4r(y + n)$
$\beta = 1/2$	$4y^2 - 4sy + (n+2)^2 - 8$	$2sy - n(n-2)$	$r(2y + n - 2)$
$\beta = 2$	$y^2 - 4sy + 4((n-1)^2 - 2)$	$2sy - 4n(n+1)$	$r(2y + 4(n+1))$

in terms of

$$r = pq, \quad s = n - 2p - 2q. \quad (6.1.3)$$

Theorem 6.2 For $\beta = 1$, we have

$$\begin{aligned} & (c_{n,q,p}^{(1)})^{-1} \int_{Gr(p, \mathbb{C}^n)} e^{x \operatorname{Tr}(I + Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-(q-p)} d\mu(Z) \\ &= \prod_1^p \frac{(n-j)!}{(q-j)!} \sum_{\ell \geq p(n-p)} \frac{p^\ell x^{\ell-p(n-p)}}{\ell!} E^{\ell,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{\substack{(i,j) \in \lambda \\ n-q < j \leq n-p}} h_{(i,j)}^\lambda \right), \\ &= \sum_{r \geq 0} x^r \sum_{\kappa \in \mathbb{Y}_r} \frac{1}{(h^\kappa)^2} \frac{(p)_\kappa (q)_\kappa}{(n)_\kappa} \\ &= {}_2F_1^{(1)}(p, q; n; y) \Big|_{\sum_\ell y_\ell^i = \delta_{1i} x} \\ &= \exp \int_0^x \frac{u(y) - p(n-p) + py}{y} dy \end{aligned} \quad (6.1.4)$$

where $u(x)$ is the unique solution to the initial value problem:

$$\left\{ \begin{array}{l} x^2 u''' + xu'' + 6xu'^2 - 4uu' + 4Qu' - 2Q'u + 2R = 0 \\ \text{with } u(x) = p(n-p) - \frac{p(n-q)}{n}x + \dots + a_{n+1}x^{n+1} + O(x^{n+1}) + \dots, \text{ near } x = 0. \end{array} \right. \quad \text{(Painlevé V)} \quad (6.1.5)$$

with a_{n+1} specified (see remark). Q and R are polynomials in x :

$$\begin{aligned} 4Q &= -x^2 + 2(n + 2(p - q))x - (n - 2p)^2 \\ 2R &= p(p - q)(x + n - 2p). \end{aligned} \quad (6.1.6)$$

The third order equation (6.1.5) has a first integral, which is second order in u and quadratic in u'' ,

$$u''^2 + \frac{4}{x^2} \left((xu'^2 + Qu' + R)u' - (u'^2 + Q'u' + R')u + \frac{1}{2}Q''u^2 - \frac{p^2(q-p)^2}{4} \right) = 0 \quad (6.1.7)$$

Remark: Note that the Painlevé equation (6.1.5) admits a solution

$$u(x) = p(n-p) - \frac{p(n-q)}{n}x + \sum_{i \geq 2} a_i x^i \quad (6.1.8)$$

with the a_i given by the indicial equation

$$i(i-1-n)(i-1+n)a_i = g_i(a_0, \dots, a_{i-1}), \quad i = 1, 2, \dots,$$

showing the existence of a free parameter at $i = n+1$. However the fact that, according to (4.2.3),

$$u(x) = p(n-p) - px + x \frac{d}{dx} \log \sum_{r \geq 0} x^r \sum_{\substack{\kappa \in \mathbb{Y}_r \\ \kappa_1^\top \leq p}} \frac{1}{(h^\kappa)^2} \prod_1^p \frac{(p-i+1)_{\kappa_i} (q-i+1)_{\kappa_i}}{(n-i+1)_{\kappa_i}}$$

leads to an explicitly known value for a_{n+1} .

Proof of Theorem 6.1: Define

$$\tilde{I}_p(t) := \int_{[-1,1]^p} |\Delta_p(y)|^{2\beta} \prod_1^p e^{\sum_{i=1}^\infty t_i y_k^i} (1 - y_k)^a (1 + y_k)^b dy_k, \quad (6.1.9)$$

and the locus

$$\mathcal{L} := \{t_1 = x \neq 0, \text{ all other } t_i = 0\}. \quad (6.1.10)$$

Using (1.0.3), and setting $a = \beta(n - q - p + 1) - 1$, $b = \beta(q - p + 1) - 1$ in (6.1.9), the linear change of variables $y_k := 2z_k - 1$ leads to:

$$\begin{aligned} \tilde{I}_p(t) \Big|_{\mathcal{L}} &= \int_{[-1,1]^p} |\Delta_p(y)|^{2\beta} \prod_1^p e^{xy_k} (1 - y_k)^{\beta(n-q-p+1)-1} (1 + y_k)^{\beta(q-p+1)-1} dy_k \\ &= c_p^{(\beta)} e^{-px} \int_{[0,1]^p} |\Delta_p(z)|^{2\beta} \prod_1^p e^{2xz_k} z_k^{\beta(q-p+1)-1} (1 - z_k)^{\beta(n-q-p+1)-1} dz_k \\ &= c_p^{(\beta)} e^{-px} \int_{Gr(p, \mathbb{F}^n)} e^{2x \operatorname{Tr}(I+Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-\beta(q-p)} d\mu(Z) \\ &= c_p^{(\beta)} e^{-px} I_p(x), \end{aligned} \quad (6.1.11)$$

with

$$c_p^{(\beta)} = 2^{p(\beta n - \beta p + \beta - 1)}. \quad (6.1.12)$$

According to the appendix, the integral $\tilde{I}_p(t)$ satisfies the Virasoro constraints (8.0.10), with $a = \beta(n - q - p + 1) - 1$, $b = \beta(q - p + 1) - 1$, and thus

$$b_0 = a - b = \beta(n - 2q), \quad b_1 = a + b = \beta(n - 2p + 2) - 2,$$

and

$$\sigma_1 = \beta n, \quad \sigma_2 = \beta(n - 1) + 1.$$

These expressions and their first t_1 - and t_2 - derivatives, evaluated along the locus $\mathcal{L}$ read as follows: ($F(t) := F_p(t) := \log \tilde{I}_p(t)$)

$$\begin{aligned} 0 &= \frac{\mathcal{J}_{-1}^{(2)} \tilde{I}_p}{\tilde{I}_p} \Big|_{\mathcal{L}} = \left(t_1 \frac{\partial}{\partial t_2} + \sigma_1 \frac{\partial}{\partial t_1} \right) F_p + p(b_0 - t_1) \Big|_{\mathcal{L}} \\ 0 &= \frac{\mathcal{J}_0^{(2)} \tilde{I}_p}{\tilde{I}_p} \Big|_{\mathcal{L}} = \left(t_1 \frac{\partial}{\partial t_3} + \sigma_2 \frac{\partial}{\partial t_2} + (b_0 - t_1) \frac{\partial}{\partial t_1} + \beta \frac{\partial^2}{\partial t_1^2} \right) F_p \\ &\quad + \beta \left(\frac{\partial F_p}{\partial t_1} \right)^2 - \frac{p}{2} (\sigma_1 - b_1) \Big|_{\mathcal{L}} \\ 0 &= \frac{\partial}{\partial t_1} \frac{\mathcal{J}_{-1}^{(2)} \tilde{I}_p}{\tilde{I}_p} \Big|_{\mathcal{L}} = \left(t_1 \frac{\partial^2}{\partial t_2 \partial t_1} + \frac{\partial}{\partial t_2} + \sigma_1 \frac{\partial^2}{\partial t_1^2} \right) F_p \Big|_{\mathcal{L}} - p \end{aligned}$$

$$\begin{aligned}
0 &= \left. \frac{\partial}{\partial t_1} \frac{\mathcal{J}_0^{(2)} \tilde{I}_p}{\tilde{I}_p} \right|_{\mathcal{L}} = \left(t_1 \frac{\partial^2}{\partial t_3 \partial t_1} + \sigma_2 \frac{\partial^2}{\partial t_2 \partial t_1} + (b_0 - t_1) \frac{\partial^2}{\partial t_1^2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_1} \right. \\
&\quad \left. + \beta \frac{\partial^3}{\partial t_1^3} \right) F_p + 2\beta \left. \frac{\partial F_p}{\partial t_1} \frac{\partial^2 F_p}{\partial t_1^2} \right|_{\mathcal{L}} \\
0 &= \left. \frac{\partial}{\partial t_2} \frac{\mathcal{J}_{-1}^{(2)} \tilde{I}_p}{\tilde{I}_p} \right|_{\mathcal{L}} = \left(t_1 \frac{\partial^2}{\partial t_2^2} + \sigma_1 \frac{\partial^2}{\partial t_1 \partial t_2} + 2 \left(\frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_1} \right) \right) F_p \Big|_{\mathcal{L}}.
\end{aligned}$$

The five equations above form a (triangular) linear system in five unknowns

$$\left. \frac{\partial F_p}{\partial t_2} \right|_{\mathcal{L}}, \quad \left. \frac{\partial F_p}{\partial t_3} \right|_{\mathcal{L}}, \quad \left. \frac{\partial^2 F_p}{\partial t_1 \partial t_2} \right|_{\mathcal{L}}, \quad \left. \frac{\partial^2 F_p}{\partial t_1 \partial t_3} \right|_{\mathcal{L}}, \quad \left. \frac{\partial^2 F_p}{\partial t_2^2} \right|_{\mathcal{L}},$$

which can be expressed in terms of

$$\frac{\partial F_p}{\partial t_1}, \quad \frac{\partial^2 F_p}{\partial t_1^2}, \quad \frac{\partial^3 F_p}{\partial t_1^3}.$$

Setting $t_1 = x$ and $F'_n = \partial F_n / \partial x$, these expressions are

$$\begin{aligned}
\left. \frac{\partial F_n}{\partial t_2} \right|_{\mathcal{L}} &= -\frac{1}{t_1} (\sigma_1 F'_n - p(t_1 - b_0)) \\
\left. \frac{\partial F_n}{\partial t_3} \right|_{\mathcal{L}} &= \frac{1}{t_1^2} \left(-2\beta t_1 (F''_n + F_n'^2) + 2(t_1^2 - b_0 t_1 + \sigma_1 \sigma_2) F'_n \right. \\
&\quad \left. + p((\sigma_1 - 2\sigma_2 - b_1)t_1 + 2b_0 \sigma_2) \right) \\
\left. \frac{\partial^2 F_n}{\partial t_1 \partial t_2} \right|_{\mathcal{L}} &= \frac{1}{t_1^2} (-\sigma_1 t_1 F''_n + \sigma_1 F'_n + b_0 p) \\
\left. \frac{\partial^2 F_n}{\partial t_1 \partial t_3} \right|_{\mathcal{L}} &= \frac{1}{2t_1^3} \left(-2\beta t_1^2 (F_n'''' + 2F'_n F''_n - F_n'^2) + 2t_1(t_1^2 - b_0 t_1 + \sigma_1 \sigma_2 + \beta) F''_n \right. \\
&\quad \left. + 2(b_0 t_1 - 2\sigma_1 \sigma_2) F'_n + p((2\sigma_2 - \sigma_1 + b_1)t_1 - 4\sigma_2 b_0) \right) \\
\left. \frac{\partial^2 F_n}{\partial t_2^2} \right|_{\mathcal{L}} &= \frac{1}{t_1^3} \left((\sigma_1^2 + 2\beta) t_1 F''_n + 2\beta t_1 F_n'^2 + (2b_0 t_1 - 2\sigma_1 \sigma_2 - \sigma_1^2) F'_n \right. \\
&\quad \left. + p((- \sigma_1 + 2\sigma_2 + b_1)t_1 - b_0(\sigma_1 + 2\sigma_2)) \right).
\end{aligned} \tag{6.1.13}$$

From (8.0.2), it follows that

$$\tilde{I}_p(t) = \begin{cases} p! \tau_p(t) & p \text{ even, } \beta = 1/2 \\ p! \tau_p(t) & p \text{ arbitrary, } \beta = 1 \\ p! \tau_{2p}(t/2) & p \text{ arbitrary, } \beta = 2, \end{cases} \quad (6.1.14)$$

where in all three cases $\tau_p(t)$ is a τ -function satisfying the KP and Pfaff-KP equations (8.0.4). Substituting (6.1.14) in the equation (8.0.4) and evaluating along the locus $\mathcal{L}$ leads to the following equations:

$$\begin{aligned} \left(\left(\frac{\partial}{\partial t_1} \right)^4 + 3 \left(\frac{\partial}{\partial t_2} \right)^2 - 4 \frac{\partial^2}{\partial t_1 \partial t_3} \right) F_p + 6 \left(\frac{\partial^2}{\partial t_1^2} F_p \right)^2 \\ = 12 \frac{p(p-1)}{(p+1)(p+2)} \frac{\tilde{I}_{p-2} \tilde{I}_{p+2}}{\tilde{I}_p^2} (1 - \delta_1^\beta) \\ \text{for } \beta = 1/2, 1, \\ \left(\left(\frac{\partial}{\partial t_1} \right)^4 + \frac{3}{4} \left(\frac{\partial}{\partial t_2} \right)^2 - \frac{\partial^2}{\partial t_1 \partial t_3} \right) F_p + 6 \left(\frac{\partial^2}{\partial t_1^2} F_p \right)^2 = \frac{3}{4} \frac{p}{p+1} \frac{\tilde{I}_{p-1} \tilde{I}_{p+1}}{\tilde{I}_p^2} \\ \text{for } \beta = 2. \end{aligned} \quad (6.1.15)$$

Then, using $F(t) = \log \tilde{I}_p(t)$, with $\tilde{I}_p$ as in (6.1.9) and (6.1.11),

$$\begin{aligned} H(y) &:= \frac{d}{dy} \log \int_{Gr(p, \mathbb{F}^n)} e^{y \operatorname{Tr}(I + Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-\beta(q-p)} d\mu(Z) \\ &= \frac{d}{dy} \log(e^{py/2} \\ &\quad \int_{[-1,1]^p} |\Delta_p(z)|^{2\beta} \prod_1^p e^{yz_k/2} (1 - z_k)^{\beta(n-q-p+1)-1} (1 + z_k)^{\beta(q-p+1)-1} dz_k) \\ &= \frac{d}{dy} F_p\left(\frac{y}{2}, 0, 0, \dots\right) + \frac{p}{2} \\ &= \frac{1}{2} \left(\frac{\partial F_p}{\partial t_1}\left(\frac{y}{2}, 0, 0, \dots\right) + p \right) \end{aligned} \quad (6.1.16)$$

satisfies the differential equation, upon substituting the derivative (6.1.13) into (6.1.15) ($H' := \frac{dH}{dy}$):

$$4 \left(y^3 H''' + 6y^3 H'^2 + (1 + \delta_1^\beta)(2y^2 H'' + 4y^2 H H' + y H^2) \right) - y P_0 H' + P_1 H + P_2$$

$$= \begin{cases} 0, & \text{for } \beta = 1, \\ 3 \frac{p(p-1)}{(p+1)(p+2)} y^3 \frac{\tilde{I}_{p-2} \tilde{I}_{p+2}}{\tilde{I}_p^2} \Big|_{\mathcal{L}}, & \text{for } \beta = 1/2, \\ \frac{3}{16} \frac{p}{p+1} y^3 \frac{\tilde{I}_{p-1} \tilde{I}_{p+1}}{\tilde{I}_p^2} \Big|_{\mathcal{L}}, & \text{for } \beta = 2, \end{cases}$$

where P_0, P_1, P_2 are polynomials in y , with coefficients depending on $r = pq$ and $s = n - 2p - 2q$, given by table (6.1.3).

From

$$\tilde{I}_p(t) \Big|_{\mathcal{L}} = c_p^{(\beta)} e^{-px} I_p(x) = 2^{p(\beta n - \beta p + \beta - 1)} e^{-px} I_p(x),$$

it follows that for $\beta = 1/2$ and 2,

$$\frac{\tilde{I}_{p-2} \tilde{I}_{p+2}}{\tilde{I}_p^2} \Big|_{\mathcal{L}} = 2^{-4} \frac{I_{p-2} I_{p+2}(x)}{I_p^2(x)} \quad \text{and} \quad \frac{\tilde{I}_{p-1} \tilde{I}_{p+1}}{\tilde{I}_p^2} \Big|_{\mathcal{L}} = 2^{-4} \frac{I_{p-1}(x) I_{p+1}(x)}{I_p^2(x)},$$

thus establishing (6.1.2). ■

Proof of Theorem 6.2: In particular, for $\beta = 1$, from Theorem 4.2, Theorem 6.1 and (6.1.5),

$$\begin{aligned} u(x) &:= x \frac{\partial}{\partial x} \log E_{x,p} \left(I_{\{\lambda \supseteq \mu\}}(\lambda) \prod_{\substack{(i,j) \in \lambda \\ n-q < j < n-p}} h_{(i,j)}^\lambda \right) \\ &= x \frac{\partial}{\partial x} \log \left(e^{-px} x^{p(n-p)} \int_{Gr(p, \mathbb{C}^n)} e^{x \operatorname{Tr}(I + Z^\dagger Z)^{-1}} \det(Z^\dagger Z)^{-(q-p)} d\mu(Z) \right) \\ &= -px + p(n-p) + xH(x) \end{aligned}$$

satisfies the differential equation (6.1.5). From $I_p(x) = 1 + \frac{pq}{n}x + \dots$, as in (4.2.6), it has the behavior near $x = 0$, spelled out in (6.1.5), namely

$$u(x) = p(n-p) - \frac{p(n-q)}{n}x + \dots \quad (6.1.17)$$

Equation (6.1.7) follows from Cosgrove and Scoufis ([14]) (see [3]), and the constant is obtained by setting $x = 0$ in the equation and using the Taylor expansion (6.1.16) of $u(x)$ about $x = 0$. ■

6.2 Hypergeometric functions, KP hierarchy and Painlevé equations

Theorem 6.3 *The hypergeometric function for $\beta = 1$ and $p, q, n \in \mathbb{C}$, expressed in t_i -variables,*

$$\tau(t) := {}_2F_1^{(1)}(p, q; n; u) \Big|_{\sum_{k=1}^m u_k^i = t_i}, \quad (6.2.1)$$

satisfies the Hirota bilinear relation as a function of $t_1, t_2, \dots$, namely for all $t, t' \in \mathbb{C}^\infty$,

$$\oint_{z=\infty} e^{\sum_1^\infty (t_i - t'_i) z^i} \tau(t - [z^{-1}]) \tau(t' + [z^{-1}]) dz = 0. \quad (6.2.2)$$

In particular, $\tau(t)$ satisfies the KP hierarchy¹⁰ ($k = 0, 1, 2, \dots$):

$$\left(s_{k+4} \left(\frac{\partial}{\partial t_1}, \frac{1}{2} \frac{\partial}{\partial t_2}, \frac{1}{3} \frac{\partial}{\partial t_3}, \dots \right) - \frac{1}{2} \frac{\partial^2}{\partial t_1 \partial t_{k+3}} \right) \tau \circ \tau = 0. \quad (6.2.3)$$

Proof: Using

$$1 - a = e^{-\sum_1^\infty \frac{a^i}{i}},$$

one has the following formula

$$\begin{aligned} \prod_{\substack{1 \leq j \leq p \\ 1 \leq k \leq m}} (1 - z_j u_k)^{-\beta} \Big|_{t_i := \frac{1}{i} \sum_{k=1}^m u_k^i} &= \prod_{\substack{1 \leq j \leq p \\ 1 \leq k \leq m}} e^{\beta \sum_{i=1}^\infty \frac{u_k^i z_j^i}{i}} \\ &= \prod_{1 \leq j \leq p} e^{\beta \sum_{i=1}^\infty z_j^i \frac{1}{i} \sum_{k=1}^m u_k^i} \\ &= \prod_{1 \leq j \leq p} e^{\beta \sum_{i=1}^\infty t_i z_j^i}. \end{aligned} \quad (6.2.4)$$

¹⁰Given a polynomial $p(t_1, t_2, \dots)$, define the customary Hirota symbol $p(\partial_t) f \circ g := p(\frac{\partial}{\partial y_1}, \frac{\partial}{\partial y_2}, \dots) f(t + y) g(t - y) \Big|_{y=0}$. The s_ℓ 's are the elementary Schur polynomials $e^{\sum_1^\infty t_i z^i} := \sum_{i \geq 0} s_i(t) z^i$ and for later use, set $s_\ell(\tilde{\partial}) := s_\ell(\frac{\partial}{\partial t_1}, \frac{1}{2} \frac{\partial}{\partial t_2}, \dots)$.

On the one hand, setting $a = \beta(n - q)$ and $b = \beta q$ in (2.3.14) using the new variables t_i , using the constant (0.0.20) and (6.2.4), the hypergeometric function equals an integral for integer $p \geq 1$,

$$\begin{aligned}
& c_{n,q,p}^{(\beta)} {}_2F_1^{(1/\beta)}(\beta p, \beta q, \beta n; u_1, \dots, u_m) \\
&= \int_{[0,1]^p} \prod_{\substack{1 \leq k \leq p \\ 1 \leq j \leq m}} (1 - z_k u_j)^{-\beta} |\Delta_p(z)|^{2\beta} \prod_{\ell=1}^p z_\ell^{\beta(q-p+1)-1} (1 - z_\ell)^{\beta(n-p-q+1)-1} dz_\ell \\
&= \int_{[0,1]^p} |\Delta_p(z)|^{2\beta} \prod_{\ell=1}^p e^{\beta \sum_{i=1}^\infty t_i z_\ell^i} z_\ell^{\beta(q-p+1)-1} (1 - z_\ell)^{\beta(n-p-q+1)-1} dz_\ell. \quad (6.2.5)
\end{aligned}$$

According to (8.0.4), this integral is a solution of the Hirota bilinear relation for $\beta = 1$, $q, n \in \mathbb{R}$, integer $p > 1$, such that $q - p + 1$ and $n - p - q + 1 > 0$.

On the other hand, the hypergeometric function, is also defined by (0.0.16), for $p, q, n \in \mathbb{R}$ and $\beta = 1$:

$${}_2F_1^{(1)}(p, q, n; u_1, \dots, u_m) \Big|_{t_i := \frac{1}{i} \sum_{k=1}^m u_k^i} = \sum_{\lambda \in \mathbb{Y}} \frac{(p)_\lambda (q)_\lambda}{(n)_\lambda h^\lambda} \mathbf{s}_\lambda(t) =: \sum_{\lambda \in \mathbb{Y}} c_\lambda \mathbf{s}_\lambda(t). \quad (6.2.6)$$

For integer $p \geq 1$, the integral (6.2.5) was shown to be a solution of the Hirota bilinear relations and so the coefficients

$$c_\lambda \Big|_{\text{integer } p \geq 1}$$

satisfy Plücker relations. They are homogeneous quadratic relations in a finite number of the c_λ 's for $\lambda \in \mathbb{Y}$ and they characterize the KP τ -functions; so, we have for every integer $p \geq 1$,

$$\begin{aligned}
0 &= \sum_{\lambda, \mu} \alpha_{\lambda, \mu} c_\lambda c_\mu, & \alpha_{\lambda, \mu} &= \pm 1 \\
&= \sum_{\lambda, \mu} \alpha_{\lambda, \mu} \frac{(p)_\lambda (q)_\lambda (p)_\mu (q)_\mu}{(n)_\lambda (n)_\mu} \\
&=: \frac{1}{Y(n)} \sum_i X_i(n, q) p^i, \quad (6.2.7)
\end{aligned}$$

where $X_i(n, q)$ are polynomials in n, q and $\sum_i$ is a finite sum. Setting $p =$ distinct integers p_k in sufficient number, one solves the homogeneous linear

system (6.2.7) in the $X_i(n, q)$ with coefficients p_k^i , whose determinant is a Vandermonde, therefore non-zero. This implies that $X_i(n, q) = 0$ for all $n, q \in \mathbb{R}$ and so, we also have

$$\sum_i X_i(n, q) p^i = 0, \quad \text{for all } p \in \mathbb{R},$$

implying the Plücker relations (6.2.6) for $p, q, n \in \mathbb{R}$. One can extend the argument further by analytic continuation to $p, q, n \in \mathbb{C}$. ■

Theorem 6.4 *The hypergeometric function*

$$H(x) = \frac{d}{dx} \log {}_2F_1^{(1)}(p, q; n; y) \Big|_{\sum_\ell y_\ell^i = \delta_{1i} x} \quad (6.2.8)$$

satisfies the Painlevé V equation:

$$4 \left(x^3 H''' + 6x^3 H'^2 + 2(2x^2 H'' + 4x^2 H H' + x H^2) \right) - x P_0 H' + P_1 H + P_2 = 0 \quad (6.2.9)$$

with P_0, P_1, P_2 as in (6.1.3).

Proof: For integer p, q , with $0 \leq p, q \leq n/2$, we have by (2.3.12),

$$\begin{aligned} {}_2F_1^{(1)}(p, q; n; u) \Big|_{\sum_\ell u_\ell^i = \delta_{1i} x} &= 1 + \sum_{r \geq 1} b_r x^r \\ &= \sum_{r \geq 0} x^r \sum_{\lambda \in \mathbb{V}_r} \frac{1}{(h^\lambda)^2} \frac{(p)_\lambda (q)_\lambda}{(n)_\lambda}, \quad p, q, n \in \mathbb{C} \end{aligned}$$

with b_r rational in p, q, n , and so

$$H(x) := \frac{d}{dx} \log {}_2F_1^{(1)}(p, q; n; y) \Big|_{\sum y_\ell^i = \delta_{1i} x} = \sum_{r \geq 0} c_r x^r,$$

also with c_r rational in p, q, n . Putting this expression into the left hand side of the Painlevé V equation leads to

$$\sum_{r=0}^{\infty} d_r x^r, \quad d_r = \text{rational in } p, q, n, \text{ with universal coefficients.}$$

For integer p, q, n with $n/2 \geq p, q \geq 0$, $H(x)$ satisfies the Painlevé equation by the proof of Theorem 6.1 and so $d_r = 0$. Using the same argument as in Theorem 6.3, we have $d_r = 0$ for all $p, q, n \in \mathbb{C}$, whenever the series ${}_2F_1^{(1)}$ makes sense. ■

7 Differential equations for the Wishart and canonical correlations distributions

Remember from section 1, the tangent space to the Grassmannians at $Id = \begin{pmatrix} I_p \\ O \end{pmatrix}$ is given by

$$T_{Id}Gr(p, \mathbb{F}^n) = \{Z \mid \text{arbitrary } (n-p) \times p \text{ matrix with values in } \mathbb{F}\}.$$

The subgroup

$$K = \left\{ B = \begin{pmatrix} B_{11} & O \\ O & B_{22} \end{pmatrix}, B_{11} \in K_1, B_{22} \in K_2 \right\}$$

acts on $T_{Id}Gr(m, \mathbb{F}^n)$ as

$$Ad_B(Z) = B_{22}ZB_{11}^{-1},$$

for which the spectrum $(\lambda_1, \dots, \lambda_m)$ of $Z^\dagger Z$ is a invariant under the action of K and $\lambda_i \geq 0$, since the matrix $Z^\dagger Z$ is positive definite.

Given $E \subseteq [0, \infty)$, consider the following probability:

$$\begin{aligned} P_{n,p}^{(\beta)}(E) &:= P(\text{all } \lambda_i \in E) \\ &= c_{p,n} \int_{\substack{Z \in T_{Id}Gr(p, \mathbb{F}^n) \\ \text{spectrum } (Z^\dagger Z) \subset E}} e^{-\text{Tr } V(Z^\dagger Z)} d\mu(Z) \\ &= \frac{\int_{E^p} |\Delta_p(z)|^{2\beta} \prod_{i=1}^n e^{-V(z_i)} z_i^{\beta(n-2p+1)-1} dz_i}{\int_{F^p} |\Delta_p(z)|^{2\beta} \prod_{i=1}^n e^{-V(z_i)} z_i^{\beta(n-2p+1)-1} dz_i}, \end{aligned} \quad (7.0.1)$$

where $d\mu(Z)$ is Haar measure (1.0.11) on $T_{Id}Gr(p, \mathbb{F}^n)$ and where $\beta = 1/2, 1, 2$ correspond to $\mathbb{R}, \mathbb{C}, \mathbb{H}$ respectively. The function $e^{-V(z)}$ will be either

- (i) Laguerre: $e^{-V(z)} = e^{-bz} z^a$, $F = \mathbb{R}^+$, $\beta = 1/2, 1, 2$
- (ii) Jacobi: $e^{-V(z)} = (1-z)^a z^b$, $F = [0, 1]$, $\beta = 1/2, 1, 2$
- (iii) Gaussian: $e^{-V(z)} = e^{-z^2}$, $F = \mathbb{R}^+$, $\beta = 1$, $n = 2p$

We shall only consider (i) and (ii), the Gaussian distribution being as in [2].

7.1 Laguerre probability on the tangent space to $Gr(p, \mathbb{F}^n)$ and the Wishart distribution

Theorem 7.1 For $e^{-V(y)} := e^{-by}y^a$, the probability (7.0.1) defined on $Z \in$ the tangent space (at the identity) to the symmetric space $Gr(p, \mathbb{F}^n)$, leads to the following probability on the (positive) spectrum $(\lambda_1, \dots, \lambda_p)$ of $Z^\dagger Z$,

$$\begin{aligned} P_{n,p}^{(\beta)}(\max_i \lambda_i \leq x) &:= c_{p,n} \int_{\substack{Z \in T_{Id} Gr(p, \mathbb{F}^n) \\ \text{spectrum}(Z^\dagger Z) \subset [0, x]}} e^{-b \text{Tr } Z^\dagger Z} (\det Z^\dagger Z)^a d\mu(Z) \\ &= \frac{\int_{[0, x]^p} |\Delta_p(y)|^{2\beta} \prod_1^p e^{-by_i} y_i^{a+\beta(n-2p)+\beta-1} dy_i}{\int_{[0, \infty)^p} |\Delta_p(y)|^{2\beta} \prod_1^p e^{-by_i} y_i^{a+\beta(n-2p)+\beta-1} dy_i} \quad (7.1.1) \end{aligned}$$

Then

$$f(x) = x \frac{d}{dx} \log P_{p,n}^{(\beta)}(\max_i \lambda_i \leq x)$$

satisfies

- for $\beta = 1$:

$$x^2 f''' + x f'' + 6x f'^2 - 4f f' - (2h f' - h' f) = 0 \quad (\text{Painlevé V})$$

with

$$2h = (a + n - 2p - bx)^2 - 4nbx.$$

- for $\beta = \frac{1}{2}$:

$$\begin{aligned} Q \left(\frac{P_{n-4, p-2}^{(\beta)} P_{n+4, p+2}^{(\beta)}}{(P_{n,p}^{(\beta)})^2} - 1 \right) - \left(3f + \frac{b^2 x^2}{2\beta} - bQ_0 x - 3Q_1 \right) f \\ = x^3 f''' - x^2 f'' + 6x^2 f'^2 - x \left(8f + \frac{b^2 x^2}{2\beta} - 2bQ_0 x - Q_2 \right) f', \quad (7.1.2) \end{aligned}$$

where

$$\alpha := a + \beta(n - 2p) + \beta - 1,$$

and

$$\begin{aligned}
Q &= \begin{cases} \frac{3}{4}p(p-1)(p+2\alpha)(p+2\alpha+1), & \text{for } \beta = 1/2 \\ \frac{3}{2}p(2p+1)(2p+\alpha)(2p+\alpha-1), & \text{for } \beta = 2 \end{cases} \\
Q_2 &= 6\beta p^2 - \frac{\alpha^2}{2\beta} + 6\alpha p + 4(1-\beta)\alpha \\
Q_1 &= 2\beta p^2 + 2\alpha p + (1-\beta)\alpha, \quad Q_0 = p + \frac{\alpha}{2\beta}.
\end{aligned} \tag{7.1.3}$$

Corollary 7.2 Consider a matrix A , which is Wishart $W_p(n, \frac{1}{2b}I_p)$ -distributed with eigenvalues $\lambda_1, \dots, \lambda_p$ (see Muirhead, p. 107). Then

$$P_{n,p}^W(x) := P_{n,p}^W(\max_i \lambda_i \leq x), \quad f := x \frac{d}{dx} \log P_{n,p}^W(x)$$

satisfy the equation

$$\begin{aligned}
&\frac{3}{4}p(p-1)n(n-1) \left(\frac{P_{n+2,p+2}^W P_{n-2,p-2}^W}{(P_{n,p}^W)^2} - 1 \right) - (3f + b^2 x^2 - bxQ_0 - 3Q_1)f \\
&= x^3 f''' - x^2 f'' + 6x^2 f'^2 - x(8f + (bx - Q_0)^2 - 4Q_1)f',
\end{aligned}$$

with

$$Q_0 = \frac{1}{2}(n+p-1) \quad \text{and} \quad 4Q_1 = (n-1)(4p+1) - p, \quad .$$

Proof: According to (5.1.2), the Wishart distribution $W_p(n, \frac{1}{2b}I_p)$ of the $p \times p$ matrix A is given by

$$P_{n,p}^W(dA) = \Gamma_p(n/2)^{-1} b^{np/2} e^{-b \operatorname{Tr} A} (\det A)^{\frac{1}{2}(n-p-1)} \prod_{1 \leq i \leq j \leq p} dA_{ij}. \tag{7.1.4}$$

and so the joint probability $P_{n,p}^W(\max_i \lambda_i \leq x)$ is precisely formula (7.1.1), with $a = p/2$ and $\beta = 1/2$,

$$\begin{aligned}
P_{n,p}^W(\max_i \lambda_i \leq x) &= c_{n,p,b} \int_{[0,x]^p} |\Delta_p(y)| \prod_{i=1}^p e^{-by_i} y_i^{\frac{1}{2}(n-p-1)} dy_i \\
&= P_{n,p}^{(\beta)}(\max_i \lambda_i \leq x) \Big|_{\substack{a=p/2 \\ \beta=1/2}}.
\end{aligned}$$

Therefore $P_{n,p}^W(x)$ also satisfies the inductive differential equation (7.1.2); we only need to check that

$$\begin{aligned}
P_{n\pm 2, p\pm 2}^W(x) &= c \int_{[0,x]^{p\pm 2}} |\Delta_{p\pm 2}(y)| \prod_1^{p\pm 2} e^{-by_i} y_i^{\frac{1}{2}(n-p-1)} dy_i \\
&= c \int_{[0,x]^{p\pm 2}} |\Delta_{p\pm 2}(y)| \prod_1^{p\pm 2} e^{-by_i} y_i^{p/2} y_i^{\frac{1}{2}((n\pm 4)-2(p\pm 2))-\frac{1}{2}} \\
&= P_{n\pm 4, p\pm 2}^{(1/2)}(\max_i \lambda_i \leq x) \text{ for } a = p/2.
\end{aligned}$$

Q, Q_0, Q_1, Q_2 can immediately be computed by setting $\beta = 1/2$, $\delta^\beta = 1$ and $2\alpha = n - p - 1$ in (7.1.3). $\blacksquare$

7.2 Jacobi probability on the tangent space to $Gr(p, \mathbb{F}^n)$ and the sample canonical correlation distribution

Theorem 7.3 For $e^{-V(y)} := (1-y)^a y^b$, the probability (7.0.1) defined on $Z \in$ the tangent space (at the identity) to the symmetric space $Gr(p, \mathbb{F}^n)$, leads to the following probability on the (positive) spectrum of $Z^\dagger Z$,

$$\begin{aligned}
P_{n,p}^{(\beta)}(\max_i \lambda_i \leq \frac{x+1}{2}) &:= c_{p,n} \int_{\substack{Z \in T_{Id} Gr(p, \mathbb{F}^n) \\ \text{spectrum}(Z^\dagger Z) \subset [0, \frac{x+1}{2}]}} e^{-\text{Tr } V(Z^\dagger Z)} d\mu(Z) \\
&= \frac{\int_{[0, \frac{x+1}{2}]^p} |\Delta_p(y)|^{2\beta} \prod_{i=1}^p (1-y_i)^a y_i^{b+\beta(n-2p+1)-1} dy_i}{\int_{[0,1]^p} |\Delta_p(y)|^{2\beta} \prod_{i=1}^p (1-y_i)^a y_i^{b+\beta(n-2p+1)-1} dy_i};
\end{aligned} \tag{7.2.1}$$

it satisfies the following differential equations, upon setting

$$f(x) := (1-x^2) \frac{d}{dx} \log P_{n,p}^{(\beta)}(\max_i \lambda_i \leq \frac{x+1}{2}),$$

- for $\beta = 1$:

$$\begin{aligned}
2(x^2 - 1)^2 f''' + 4(x^2 - 1) (x f'' - 3f'^2) + (16xf - u(x^2 - 1) - 2sx - r) f' \\
- f(4f - ux - s) = 0
\end{aligned} \tag{7.2.2}$$

$$\begin{aligned}
& \bullet \text{ for } \beta = \left\{ \frac{1}{2} : \right. \\
& \quad \left. Q \left(\frac{P_{n+\frac{1}{2}, p+\frac{1}{2}}^{(\beta)} P_{n-\frac{1}{2}, p-\frac{1}{2}}^{(\beta)}}{(P_{n,p}^{(\beta)})^2} - 1 \right) \right. \\
& = 4(u+1)(x^2-1)^2 \left(-u(x^2-1)f''' + (12f - ux - 3s)f'' + 6u(u-1)f'^2 \right) \\
& \quad - (x^2-1)f' \left(24f(u+3)(2f-s) + 8fu(5u-1)x - u(u+1)(ux^2 + 2sx + 8) + Q_2 \right) \\
& \quad + f \left(48f^3 + 48f^2(ux + 2x - s) + 2f(8u^2x^2 + 2ux^2 - 12usx - 24sx + Q_4) \right. \\
& \quad \left. - u(u+1)x(3ux^2 + sx - 2ux - 3u) + Q_3x - Q_1s \right), \tag{7.2.3}
\end{aligned}$$

where

$$b_0 = a - b - \beta(n - 2p + 1) + 1, \quad b_1 = a + b + \beta(n - 2p + 1) - 1$$

$$\begin{aligned}
r &= \frac{2}{\beta}(b_0^2 + (b_1 + 2 - 2\beta)^2) & s &= \frac{2}{\beta}b_0(b_1 + 2 - 2\beta) \\
u &= \frac{2}{\beta}(2\beta p + b_1 + 2 - 2\beta)(2\beta p + b_1),
\end{aligned}$$

and

$$\begin{aligned}
Q &= \frac{3}{16}((s^2 - ur + u^2)^2 - 4(rs^2 - 4us^2 - 4s^2 + u^2r)) \\
Q_1 &= 3s^2 - 3ur - 6r + 2u^2 + 23u + 24 \\
Q_2 &= 3us^2 + 9s^2 - 4u^2r + 2ur + 4u^3 + 10u^2 \\
Q_3 &= 3us^2 + 6s^2 - 3u^2r + u^3 + 4u^2 \\
Q_4 &= 9s^2 - 3ur - 6r + u^2 + 22u + 24 = Q_1 + (6s^2 - u^2 - u). \tag{7.2.4}
\end{aligned}$$

Corollary 7.4 *Let $A = ZZ^\top$ have the Wishart distribution $W_{p+q}(n, \Sigma)$ -distribution. Break up the matrices Σ and A , as follows:*

$$\begin{array}{ccc}
& \xleftrightarrow{p} & \xleftrightarrow{q} \\
\Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{12}^\top & \Sigma_{22} \end{pmatrix} & \begin{array}{c} \downarrow p \\ \downarrow q \end{array} & A = \begin{pmatrix} A_{11} & A_{12} \\ A_{12}^\top & A_{22} \end{pmatrix} \begin{array}{c} \downarrow p \\ \downarrow q \end{array}
\end{array}$$

Assume the eigenvalues $\rho_1^2, \dots, \rho_p^2$ of $\Sigma_{11}^{-1} \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{12}^\top$ all zero. Then the probability distribution of the eigenvalues $r_1^2, \dots, r_p^2$ of $A_{11}^{-1} A_{12} A_{22}^{-1} A_{12}^\top$ (sample canonical correlation coefficients) is given by

$$\begin{aligned} P_{n,p,q}^C(x) &= P(0 \leq r_i^2 \leq \frac{1+x}{2} \text{ for } 1 \leq i \leq p) \\ &= \frac{\int_{[0, \frac{1+x}{2}]^p} |\Delta_p(z)| \prod_{i=1}^p z_i^{\frac{1}{2}(q-p-1)} (1-z_i)^{\frac{1}{2}(n-p-q-1)} dz_i}{\int_{[0,1]^p} |\Delta_p(z)| \prod_{i=1}^p z_i^{\frac{1}{2}(q-p-1)} (1-z_i)^{\frac{1}{2}(n-p-q-1)} dz_i} \end{aligned}$$

and

$$f(x) := (1-x^2) \frac{d}{dx} \log P_{n,p,q}^C(x)$$

satisfy the inductive PDE:

$$Q \left(\frac{P_{n+4,p+2,q+2}^C P_{n-4,p-2,q-2}^C}{P_{n,p,q}^C{}^2} - 1 \right) = \left\{ \begin{array}{l} \text{same expression as the} \\ \text{right hand side of (7.2.3)} \end{array} \right\},$$

with Q_1, Q_2, Q_3 being symmetric polynomials of p, q , given by (7.2.4), where

$$\begin{aligned} u &= n(n-2) \\ r/4 &= n^2/2 - n(p+q) + p^2 + q^2 \\ s &= (n-2p)(n-2q) \\ Q &= 48pq(p-1)(q-1)(n-p)(n-q)(n-p-1)(n-q-1). \end{aligned}$$

Remark: For instance

$$\begin{aligned} Q_1 &= 24(n-1)(p^2+q^2-n(p+q)) + 48pq(n^2-n(p+q)+pq) \\ &\quad - (n-6)(n-1)^2(n+4). \end{aligned}$$

Proof: The proof follows immediately from Corollary 5.2 and Theorem 7.3. ■

8 Appendix: The Pfaff-KP hierarchy and Virasoro constraints

Consider weights of the form $\rho(z)dz := e^{-V(z)}dz$ on an interval $F = [A, B] \subseteq \mathbb{R}$, with rational logarithmic derivative and subjected to the following bound-

ary conditions:

$$-\frac{\rho'}{\rho} = V' = \frac{g}{f} = \frac{\sum_0^\infty b_i z^i}{\sum_0^\infty a_i z^i}, \quad \lim_{z \rightarrow A, B} f(z) \rho(z) z^k = 0 \text{ for all } k \geq 0, \quad (8.0.1)$$

Theorem 8.1 *The multiple integrals*

$$\begin{aligned} I_p(t; \beta) &= \int_{F^p} |\Delta_p(z)|^{2\beta} \prod_{k=1}^p \left(e^{\sum_1^\infty t_i z_k^i} \rho(z_k) dz_k \right), \text{ for } p > 0 \\ &= \begin{cases} p! \tau_p(t) & p \text{ even}, & \beta = 1/2 \\ p! \tau_p(t) & p \text{ arbitrary}, & \beta = 1 \\ p! \tau_{2p}(t/2) & p \text{ arbitrary}, & \beta = 2 \end{cases} \end{aligned} \quad (8.0.2)$$

with $I_0 = 1$, satisfy

(i) the following Virasoro constraints for all $k \geq -1$:

$$\sum_{i \geq 0} \left(a_i {}^\beta \mathbb{J}_{k+i,p}^{(2)}(t, p) - b_i {}^\beta \mathbb{J}_{k+i+1,p}^{(1)}(t, p) \right) I_p(t; \beta) = 0, \quad (8.0.3)$$

in terms of the coefficients a_i, b_i of the rational function $(-\log \rho)'$

(ii) The Pfaff-KP hierarchy: (see footnote 14 for notation)

$$\left(\mathbf{s}_{k+4}(\tilde{\partial}) - \frac{1}{2} \frac{\partial^2}{\partial t_1 \partial t_{k+3}} \right) \tau_p \circ \tau_p = (1 - \delta_1^\beta) \mathbf{s}_k(\tilde{\partial}) \tau_{p+2} \circ \tau_{p-2} \quad (8.0.4)$$

$p \text{ even}, k = 0, 1, 2, \dots$

of which the first equation reads ($p \text{ even}$)

$$\begin{aligned} &\left(\left(\frac{\partial}{\partial t_1} \right)^4 + 3 \left(\frac{\partial}{\partial t_2} \right)^2 - 4 \frac{\partial^2}{\partial t_1 \partial t_3} \right) \log \tau_p + 6 \left(\frac{\partial^2}{\partial t_1^2} \log \tau_p \right)^2 \\ &= 12 \frac{\tau_{p-2} \tau_{p+2}}{\tau_p^2} (1 - \delta_{\beta,1}). \end{aligned} \quad (8.0.5)$$

(iii) More generally, the functions $\tau(t)$ satisfy the Hirota bilinear relations for all $t, t' \in \mathbb{C}^\infty$ and m, p positive integers (see footnote 10 for notation)

• $\beta = 1$

$$\oint_{z=\infty} \tau_p(t - [z^{-1}]) \tau_p(t' + [z^{-1}]) e^{\sum_1^\infty (t_i - t'_i) z^i} dz = 0, \quad (8.0.6)$$

- $\beta = 1/2$ and 2

$$\oint_{z=\infty} \tau_{2p}(t - [z^{-1}]) \tau_{2m+2}(t' + [z^{-1}]) e^{\sum_0^\infty (t_i - t'_i) z^i} z^{2p-2m-2} dz \\ + \oint_{z=0} \tau_{2p+2}(t + [z]) \tau_{2m}(t' - [z]) e^{\sum_0^\infty (t'_i - t_i) z^{-i}} z^{2p-2m} dz = 0, \quad (8.0.7)$$

Example (Jacobi β -integral)

This case is particularly important, because it covers the integrals in Theorems 0.2 and 0.3. The weight and the a_i and b_i , as in (8.0.1), are given by

$$\rho(z) := e^{-V} = (1-z)^a (1+z)^b, \quad V' = \frac{g}{f} = \frac{a-b+(a+b)z}{1-z^2}$$

$$a_0 = 1, a_1 = 0, a_2 = -1, b_0 = a-b, b_1 = a+b, \text{ and all other } a_i, b_i = 0.$$

The integrals

$$I_p = \int_{E^p} |\Delta_p(z)|^{2\beta} \prod_{k=1}^p (1-z_k)^a (1+z_k)^b e^{\sum_{i=1}^\infty t_i z_k^i} dz_k \quad (8.0.8)$$

satisfy the Virasoro constraints ($k \geq -1$):

$$\mathcal{J}_k^{(2)} I_p = \left({}^\beta \mathbb{J}_{k+2,p}^{(2)} - {}^\beta \mathbb{J}_{k,p}^{(2)} + b_0 {}^\beta \mathbb{J}_{k+1,p}^{(1)} + b_1 {}^\beta \mathbb{J}_{k+2,p}^{(1)} \right) I_p = 0. \quad (8.0.9)$$

Introducing $\sigma_i = (2p - i - 1)\beta + i + 1 + b_1$,

Then introducing the function $F_p := \log \tau_p(t)$, the two first Virasoro constraints for $m = 1, 2$ divided by τ_p are given by

$$\frac{\mathcal{J}_{-1}^{(2)} \tau_p}{\tau_p} = \left(\sum_{i \geq 1} i t_i \frac{\partial}{\partial t_{i+1}} - \sum_{i \geq 2} i t_i \frac{\partial}{\partial t_{i-1}} + \sigma_1 \frac{\partial}{\partial t_1} \right) F_p + p(b_0 - t_1) = 0 \\ \frac{\mathcal{J}_0^{(2)} \tau_p}{\tau_p} = \left(\sum_{i \geq 1} i t_i \left(\frac{\partial}{\partial t_{i+2}} - \frac{\partial}{\partial t_i} \right) + b_0 \frac{\partial}{\partial t_1} + \beta \frac{\partial^2}{\partial t_1^2} + \sigma_2 \frac{\partial}{\partial t_2} \right) F_p \\ + \beta \left(\frac{\partial F_p}{\partial t_1} \right)^2 - \frac{p}{2} (\sigma_1 - b_1) = 0. \quad (8.0.10)$$

References

- [1] M. Adler and P. van Moerbeke: *Bäcklund transformations, Birkhoff strata and Isospectral sets of differential operators*, Advances in Mathematics, **108**, 140-204, (1994).
- [2] M. Adler and P. van Moerbeke: *Hermitian, symmetric and symplectic random ensembles: PDE's for the distribution of the spectrum*, Annals of Mathematics, **153**, 149–189 (2001). (math-ph/0009001)
- [3] M. Adler and P. van Moerbeke: *Integrals over classical groups, random permutations, Toda and Toeplitz lattices*, Comm. Pure Appl. Math. **54**, 153–205 (2001) (math.CO/9912143)
- [4] M. Adler, T. Shiota and P. van Moerbeke: *Random matrices, Virasoro algebras and non-commutative KP*, Duke Math. J. **94**, 379-431 (1998).
- [5] M. Adler, T. Shiota and P. van Moerbeke: *Pfaff τ -functions* , Math. Annalen (2001) (solv-int/9909010)
- [6] D. Aldous and P. Diaconis: *Longest increasing subsequences: From patience sorting to the Baik-Deift-Johansson theorem*, Bull. Am. Math. Soc. (new series) **36** (4), 413–432 (1999).
- [7] K. Aomoto: *Jacobi polynomials associated with Selberg integrals* , SIAM J. Math. Anal. **18** , 545–549 (1987).
- [8] B. Baik, P. Deift and K. Johansson: *On the distribution of the length of the longest increasing subsequence of random permutations*, Journal Amer. Math. Soc. **12** , 1119-1178 (1999) (MathCO/9810105).
- [9] J. Baik and E. Rains: *Algebraic aspects of increasing subsequences*, math.CO/9905083 **B** , – (1999).
- [10] A. Borodin, G. Olshanski: *Z-Measures on partitions, Robinson-Schensted-Knuth correspondence, and $\beta = 2$ random matrix ensembles*, “Random matrices and their applications”, Mathematical Sciences research Institute Publications #40, Cambridge University Press, (2001) (MathCO/9905189)

- [11] A. Borodin, G. Olshanski: *Distributions on partitions, point processes, and the hypergeometric kernel* Comm. Math. Phys. **211**, 335–358 (2001). (math.RT/9904010)
- [12] A. Borodin, A. Okounkov, G. Olshanski: *Asymptotics of Plancherel measures for symmetric groups* J. Amer. Math. Soc. (to appear) (math.CO/9905032)
- [13] A.G. Constantine : *Some non-central distribution problems in multivariate analysis* , Ann. of Math. Statist. **34**, 1270-1285 (1963).
- [14] C. M. Cosgrove, G. Scoufis: *Painlevé classification of a class of differential equations of the second order and second degree* , Studies. Appl. Math. **88** , 25–87 (1993).
- [15] P. Diaconis, M. Shashahani: *On the eigenvalues of random matrices* J. Appl. Prob., suppl. in honour of Takàcs **31A**, 49-61 (1994).
- [16] E. Dueñez: *Random matrix ensembles associated to compact symmetric spaces*, Thesis, Princeton University 2001.
- [17] P.J. Forrester: *The spectrum edge of random matrix ensembles* , Nucl. Phys. B, **402**, 709-728 (1993).
- [18] P.J. Forrester: *Painlevé transcendent evaluation of the scaled distribution of the smallest eigenvalue in the Laguerre orthogonal and symplectic ensembles*,
- [19] I. M. Gessel: *Symmetric functions and P-recursiveness* , J. of Comb. Theory, Ser A, **53**, 257–285 (1990)
- [20] C. Greene: *An extension of Schensted's theorem*, Adv. in Math. **14**, 254–265 (1974)
- [21] Harish-Chandra: *Differential operators on a semi-simple Lie algebra* Amer. J. of Math., **79**, 87–120 (1957).
- [22] S. Helgason: *Groups and geomatric analysis; integral geometry, invariant differential operators, and spherical functions*, Acad. Press 1984
- [23] S. Helgason: *Differential geometry and symmetric spaces*, Acad. Press 1962

- [24] H. Hotelling: *Relations between two sets of variates*, Biometrika, **28** 321–377 (1936).
- [25] A. T. James: *Normal multivariate analysis and the orthogonal group*, Ann. Math. Statist. **25** 40–75 (1954).
- [26] I. M. Johnstone: *On the distribution of the largest principal component*, preprint, August 2000.
- [27] K. Johansson: *On random matrices from the compact classical groups*, Ann. of Math., **145**, 519–545 (1997)
- [28] K. Johansson: *The Longest Increasing Subsequence in a Random Permutation and a Unitary Random Matrix Model*, Math. Res. Lett., **5**, no. 1-2, 63–82 (1998)
- [29] K. W. J. Kadell : *The Selberg-Jack polynomials* Adv. in Math., – (199?).
- [30] J. Kaneko : *Selberg integrals and hypergeometric functions associated with Jack polynomials* SIAM J. Math. Anal. **24**, 1086–1110 (1993).
- [31] Kerov : *The boundary of Young tableaux and random Young tableaux*, DIMACS series in “Discrete Mathematics and Computer Science”, vol. **24**, 133–158 (1996).
- [32] D. Knuth: “The Art of Computer programming, Vol III: Searching and Sorting”, Addison-Wesley, Reading, MA, 1973.
- [33] B.F. Logan and L.A. Shepp: *A variational problem for random Young tableaux*, Advances in Math., **26**, 206–222 (1977).
- [34] I.G. MacDonald: “Symmetric functions and Hall polynomials”, Clarendon Press, 1995.
- [35] R.J. Muirhead: “Aspects of Multivariate Statistical Theory”, Wiley series in probability and Mathematical statistics, 1982.
- [36] A. Okounkov: *Random matrices and random permutations*, – Math.CO/99-03176, (1999).
- [37] A.Yu. Orlov, D.M. Scherbin: *Multivariate hypergeometric functions as tau functions of Toda lattice and Kadomtsev-Petviashvili equation*, math-ph/0003011 (2000).

- [38] A.Yu. Orlov, D.M. Scherbin: *Fermionic representation for basic hypergeometric functions related to Schur polynomials*, nlin.SI/0001001 (2001).
- [39] E. M. Rains: *Topics in Probability on compact Lie groups*, Harvard University doctoral dissertation, (1995).
- [40] E. M. Rains: *Increasing subsequences and the classical groups*, Elect. J. of Combinatorics, **5**, R12, (1998).
- [41] B.E. Sagan: *The Symmetry Group*, Wadsworth & Brooks, Pacific Grove, California, 1991.
- [42] P. Sarnak: *Arithmetic quantum chaos*, Israel Math. Conf. Proceedings, **8**, 183–236 (1995).
- [43] R. P. Stanley: *Some combinatorial properties of Jack symmetric functions*, Adv. in Math., **77**, 76–115 (1989).
- [44] R. P. Stanley: *Enumerative Combinatorics*, Cambridge Studies in Advanced Math. 49, Vols 1 and 2 (1997).
- [45] D. Stanton, D. White: *Constructive Combinatorics*, Springer-Verlag, NY (1986).
- [46] A. Terras: “Harmonic analysis on Symmetric Spaces and Applications II”, Springer-verlag, 1988.
- [47] C.L. Terng: *Isoparametric submanifolds and their Coxeter groups*, J. Differential Geometry. **21** 79–107 (1985), .
- [48] C.A. Tracy and H. Widom: *On the distribution of the lengths of the longest monotone subsequences in random words*, math.CO/9904042, (1999).
- [49] P. van Moerbeke: *Integrable lattices: random matrices and random permutations*, ”Random matrices and their applications”, Mathematical Sciences research Institute Publications #40, Cambridge University Press, pp 321-406, (2001)
- [50] A.M. Vershik and S.V. Kerov: *Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tables*, Soviet Math. Dokl., **18**, 527–531 (1977).